

Pre and Post Break Parameter Inference*

Graham Elliott

University of California, San Diego

9500 Gilman Drive

La Jolla, CA, 92093-0508

Ulrich K. Müller

Princeton University

Princeton, NJ, 08544-1021

Preliminary: Please Do Not Cite or Circulate

First Draft: September 2008

This Version: October 2008

Abstract

This paper discusses inference about the pre and post break value of a scalar parameter in GMM time series models with a single break at an unknown point in time. We show that treating the break date estimated by least squares as the true break date leads to substantially oversized tests and confidence intervals unless the break is large. We develop an alternative test that controls size uniformly and that is approximately efficient in some well defined sense.

JEL Classification: C32, C12

Keywords: Nuisance Parameters; Least Favorable Distribution; Size of Test; Monte Carlo; Uniformity

*The authors thank Jia Li for excellent research assistance. Müller gratefully acknowledges the NSF for financial support via grant SES-0751056.

1 Introduction

This paper is concerned with testing hypotheses about the pre and post break value of a parameter in a time series model with a single break. By reversing the time ordering, inference about the post break value becomes inference about the pre break value, so that for simplicity, we will exclusively discuss the post break case. If the break date is known, inference is straightforward, as standard results apply after restricting attention to the stable post break data. The effect of an incorrectly chosen break date crucially depends on whether the chosen date is earlier or later than the true break date. On the one hand, if the chosen date is later, then a restriction to the presumed post break data still yields a stable model, and inference remains valid. There is a cost of efficiency, though, since more post break data could have been used. On the other hand, if the chosen date is earlier than the true break date, then the presumed post break data stems from an unstable model. Parameter estimators in this unstable model tend to yield estimates of the average parameter value, which is different from the true post break value, so standard inference yields tests and confidence intervals with distorted size.

These distortionary effects are small if the chosen break date is sufficiently close to the true break date. With the break date unknown, this requires a precise break date estimator. As formally shown by Bai (1994, 1997) and Bai and Perron (1998) for linear regressions, and Hall, Han, and Boldea (2008) for two stage least squares, the least squares break date estimator is indeed sufficiently precise for these distortionary effects to become asymptotically negligible *if* the parameter shift is sufficiently pronounced. Formally, these papers study asymptotics in which the break magnitude, while possibly shrinking, is outside the local $T^{-1/2}$ neighborhood. Relative to the sampling uncertainty of the parameter estimator in a stable model, the parameter shift thus becomes infinitely large. We consider the behavior of 5% nominal level two-sided test based on the least squares break date estimator under local asymptotics in Section 2.1.2 below. With the break restricted to the middle 70% of the sample, the largest null rejection probability is almost 30%, a break magnitude of less than 5 standard deviations leads to effective size of more than 10% for all break dates, and a break of 11 standard deviations still yields size greater than 10% for some break dates. These distortions are further exacerbated by an attempt to pre-test for parameter stability. In other words, for any sample size T , there exist a break magnitude for which these standard methods yield incorrect inference, so they are not uniformly valid.

It is instructive to consider these issues in the context of an example. Productivity growth in most developed nations appears to have suffered shifts in its mean a number of times. Most researchers agree on a downward shift in productivity following the oil crisis of 1973, typically dated at the third quarter. In post 1973 data, there is a suggestion of a mid 1990's upward shift




Figure 1: Quarterly growth in non farm business productivity growth and average growth pre and post 1995 Q4 (BLS series PRS85006092)

Table 1: Inference about current US productivity growth

| Method | 95% Confidence Interval |
|------------------------------------|-------------------------|
| Break date chosen | [1.36,3.42] |
| Least squares break date estimator | [1.97,3.27] |
| Test of this paper | [1.99,3.37] |

in average productivity growth and speculation about its cause. A recent summary appears in Jorgenson, Ho, and Stiroh (2008), who informally suggest a break in the fourth quarter of 1995. Gordon (2003) also dates the increase to 1995. Anderson and Klieson (2006) refer to the increase in productivity as the 'defining economic event of the past decade'.

Figure 1 plots quarterly observations of US non farm business productivity growth, along with the pre and post break means using Jorgenson, Ho, and Stiroh (2008) break date of 1995 Q4. Based on this data, quarterly productivity growth increased by 1.1% in 1995 Q4. The least-squares estimate of the break date is slightly later at 1997 Q1. The first two rows of Table 1 contain 95% confidence intervals for current US productivity growth, conditional on these two break dates. The parameter stability test of Elliott and Müller (2006) fails to reject the null hypothesis of stable mean growth on the 10% level.

With a (long-run) standard deviation of quarterly productivity growth of approximately 3% and a sample of $T = 136$ observations, 5-11 standard deviations of the full sample estimator

correspond to a 1.3-2.8% shift in quarterly productivity growth. Absent prior knowledge that the break date is indeed 1995 Q4, or that the change of productivity is at least of the order of 1.5%, nothing suggests that either interval has close to its nominal confidence level. More generally, it seems hard to argue that break magnitudes of less than 11 standard deviations of the full sample estimator are not part of the empirically relevant parameter space in most applications. One way to see this is that shifts of, say, 8 standard deviations tend to induce highly significant rejections of the null hypothesis of parameter stability with high probability—see Elliott and Müller (2007). Yet there is continued debate about the stability of monetary policy, for instance, with Orphanides (2004) arguing for rather stable relationships, while Cogley and Sargent (2005) find instabilities that they deem ‘economically important’, but of a magnitude that would be detected by a formal parameter stability test less than 25% of the time.

Simple adjustments to the standard procedure that ensure size control over all break magnitudes do not deliver reasonable tests. For instance, in the set-up mentioned above and described in detail in Section 2.1.2, the critical value for the 5% two-sided t-statistic would need to be increased from 1.96 to approximately 4.9 to ensure uniform size control, with obvious adverse effects on power.¹ A Bonferroni procedure based on uniformly valid confidence sets for the break date developed by Elliott and Müller (2007) performs well for large breaks, but has poor power for breaks of moderate magnitude.

The main contribution of this paper is the construction of a powerful test about the post break parameter value in a general GMM time series model with unknown break date that controls size uniformly over the break magnitude. This test follows a switching rule: if there is strong evidence for a large break, then the test essentially reduces to standard inference using post break data determined by the least squares break date estimator, with a slight increase of the 5% critical value from 1.96 to 2.01. In absence of strong evidence for a large break, the test switches to a likelihood ratio test. This likelihood ratio test is carefully constructed to ensure both overall size control and approximate efficiency in the sense that for a particular weighting function, (local asymptotic) weighted average power of the suggested test is demonstrably at most 2.5% smaller than the weighted average power of any test that controls size.

From a statistical point of view, the problem is one of constructing powerful tests in the presence of a nuisance parameter under the null hypothesis—the nuisance parameter being the break data and break magnitude, neither of which can be consistently estimated under local asymptotics. We appropriately modify the algorithm developed in Müller and Watson (2008a) to identify the approximate least favorable distribution for this nuisance parameter.

¹This adjustment to the critical value is also the end result of the size corrected hybrid subsampling method recently advanced by Andrews and Guggenberger (2007a, 2007b).

This approximate least favorable distribution identifies the null density for the likelihood ratio part of the test statistic, and, using the insight of Müller and Watson (2008a) and Andrews, Moreira, and Stock (2007), allows the construction of an upper bound on the power of all tests that control size.² Beyond overcoming the substantial 'engineering' challenges in this example with a two dimensional nuisance parameter, exacerbated by the impossibility of accurately representing continuous time Gaussian processes on a discrete computer, a further contribution of this paper to this methodology is a careful consideration of size control. A test controls size if the rejection probability, viewed as a function on the null parameter space, takes on values below or at the nominal level. The standard approach is to evaluate the rejection probability for a grid of parameter points by Monte Carlo, and to conclude size is controlled if none of the estimates is larger than the nominal level. We develop an alternative algorithm that, while still based on Monte Carlo estimation, instead estimates (an upper bound on) the whole rejection probability function, ensuring that no additional implicit smoothness assumptions are necessary. It is important to note, however, that all of these 'engineering' challenges only concern the *derivation* of the suggested test; its *application* for a given sample is essentially no more difficult than the estimation of the GMM model over 142 different subsamples.

In many ways, the analysis in this paper mirrors the developments in the weak instrument literature over the last decade: Similar to Staiger and Stock (1997), we consider an alternative asymptotic embedding that formalizes the lack of uniform validity of the standard method. Akin to their rule of thumb that standard asymptotics are approximately reliable when the first stage F-statistic is sufficiently large, we suggest a formal switching rule to the least squares break date based inference when a parameter stability test is highly significant. Finally, as Andrews, Moreira and Stock (2006, 2007), we consider tests that are efficient relative to a weighted average power criterion, and identify an approximately optimal test.

Returning to the US productivity example, the last row in Table 1 shows the 95% confidence interval constructed by inverting the test suggested here. In this example, the interval is only slightly wider than the one based on the least squares based break date estimator, and remains informative about current US productivity.

The remainder of the paper is organized as follows. The next section defines the asymptotic analogues to the small sample post break parameter inference problem, quantifies the lack of size control of inference based on the least squares break date estimator, summarizes our solution and discusses the relationship of the small sample problem in parametric and GMM models to

²By reframing the issue of identifying a powerful test with uniform size control in general decision theoretic terms (see Section 3.2 below), this power bound is closely related to the bound on the minimax value generated by the least favorable distribution over a subset of possible distributions, as analyzed by Chamberlain (2000).

the limiting problem. In Section 3, we discuss in detail the construction of the suggested test, the determination of its size and the bound on power. The test statistic is defined in terms of partial sample GMM statistics (cf. Andrews (1993)) in Section 4, where we also consider its asymptotic and small sample properties. Proofs not given in the main text are collected in an appendix.

2 Limiting Problem and Underlying Models

The natural large sample analogue to small sample post break parameter inference is an inference problem about the drift component of a Gaussian process on the unit interval. The scalar version of this limiting problem is discussed first. We investigate the corresponding asymptotic properties of least squares break date based inference, and provide an overview of our line of attack, along with the key properties of the solution we provide. The second subsection considers the multivariate limiting problem, and shows in which sense the approximate efficiency of our solution in the scalar case carries over to the multivariate case. Finally, subsection 3 provides a formal link between the underlying small sample problem to the multivariate limiting problem for both parametric and GMM time series models.

2.1 Scalar Limiting Problem

2.1.1 Statement of the problem

The main focus of this paper is the following hypothesis testing problem: Suppose we observe the scalar Gaussian process $G(\cdot)$ on the unit interval (a random element with values in $D_{[0,1]}$, the space of cadlag functions on the unit interval)

$$G(s) = W(s) + \beta s + \delta \min(\rho, s), \quad s \in [0, 1] \quad (1)$$

where $W(\cdot)$ is standard Wiener process and $\beta, \delta \in \mathbb{R}$, and $\rho \in [0.15, 0.85]$. The process $G(s)$ is a Wiener process with drift $\beta + \delta$ for $s < \rho$, and drift β for $s \geq \rho$. The testing problem is $H_0 : \beta = 0$ against $H_1 : \beta \neq 0$, so that geometrically, the question is whether the slope of the deterministic component $\beta s + \delta \min(\rho, s)$ is equal to zero after the (potential) kink at $s = \rho$. The location ρ and magnitude δ of the kink are nuisance parameters. This composite nature of the hypotheses can be made more explicit by introducing the notation $\theta = (\beta, \delta, \rho)$, $\Theta_0 = \{\theta = (0, \delta, \rho) : \delta \in \mathbb{R} \text{ and } \rho \in [0.15, 0.85]\}$ and $\Theta_1 = \{\theta = (\beta, \delta, \rho) : \beta \neq 0, \delta \in \mathbb{R} \text{ and } \rho \in [0.15, 0.85]\}$, so that the hypothesis testing problem becomes

$$H_0 : \theta \in \Theta_0 \quad \text{against} \quad H_1 : \theta \in \Theta_1. \quad (2)$$

Let $\varphi : D_{[0,1]} \mapsto [0, 1]$ be a possibly randomized test of (2), where $\varphi(g)$ indicates the conditional probability of rejecting the null hypothesis test conditional on observing $G = g$. If the range of φ only consists of the two values $\{0, 1\}$, then φ is not randomized. The unconditional rejection probability is then simply the expectation of $\varphi(G)$, which we write as $E_\theta[\varphi(G)]$ to indicate that this expectation depends on the value of θ . With these definitions, a test φ is of level 5% if

$$E_\theta[\varphi(G)] \leq 5\% \text{ for all } \theta \in \Theta_0. \quad (3)$$

To illustrate the link to the post break parameter inference problem, consider observations $\{y_t\}_{t=1}^T$ from a scalar Gaussian model with a break in the mean at time τ

$$y_t = \mu + \Delta \mathbf{1}[t \leq \tau] + \varepsilon_t, \quad \varepsilon_t \sim \mathcal{N}(0, 1). \quad (4)$$

We then have the following equality in distribution

$$T^{-1/2} \sum_{t=1}^{sT} y_t \sim G(s) \quad \text{for any } s \in \left\{ \frac{t}{T} \right\}_{t=1}^T$$

where $\rho = \tau/T$, $\beta = T^{1/2}\mu$ and $\delta = T^{1/2}\Delta$. Except for the scaling by $T^{-1/2}$ and the discretization $s \in \left\{ \frac{t}{T} \right\}_{t=1}^T$, the testing problem (2) involving the observation G is therefore identical to inference about the post-break mean in Gaussian location model (4), with the break date restricted to be in the middle 70% of all observations. We argue below how the testing problem (2) based on the observation $G(\cdot)$ naturally arises as the asymptotically relevant problem in more complicated settings, but first discuss this limiting problem in more detail.

2.1.2 Tests Based on Least-Square Break Date Estimator

As discussed in the introduction, Bai (1994, 1997) and Bai and Perron (1998) suggest and analyze the following procedure for conducting inference about the post break value of a coefficient in a linear regression that is subject to a break: Estimate the break date by least-squares, construct a post break dummy using this estimate, and perform the usual t-test on that dummy. These papers show that this procedure results in asymptotically correct inference for break magnitudes that diverge when multiplied by the square root of the sample size. This corresponds to $|\delta| \rightarrow \infty$ in the notation developed here. Another procedure, arguably most prevalent in applied work, first performs a pretest for a break on some conventional significance level, followed by standard full sample inference about the post break value if no significant break is found, and performs least-squares based post break inference as above if the pretest rejects.

In the limiting problem (1), the nominally 5% level least squares and pretest procedure (based on the supF statistic) correspond to the test

$$\mathbf{1}[\text{supF} > \text{cv}_{pre}(\alpha_{pre})]\mathbf{1}[\hat{t} > \Phi(0.975)] + \mathbf{1}[\text{supF} < \text{cv}_{pre}(\alpha_{pre})]\mathbf{1}[|G(1)| > \Phi(0.975)] \quad (5)$$

where $\text{cv}_{pre}(\alpha_{pre})$ is the critical value of the supF statistic of level α_{pre} as tabulated in Andrews (1993), (and $\text{cv}_{pre}(1) = 0$ corresponds to always using the least squares break date estimator, with no pretest), $\Phi(0.975) \simeq 1.96$ is the 97.5th percentile of a standard normal and

$$\hat{\rho} = \underset{r \in [0.15; 0.85]}{\text{argmax}} \frac{(G(r) - rG(1))^2}{r(1-r)}, \quad \text{supF} = \frac{(G(\hat{\rho}) - \hat{\rho}G(1))^2}{\hat{\rho}(1-\hat{\rho})}, \quad \hat{t} = \frac{G(1) - G(\hat{\rho})}{\sqrt{1-\hat{\rho}}}. \quad (6)$$

Under standard assumptions, the large sample properties of the small sample pretest and least square based tests in a linear regression with a single coefficient converge in distribution to those in (5) for breaks of magnitude $T^{-1/2}$ —this follows, for instance, from Proposition 1 of Elliott and Müller (2007).

Figure 1 displays the null rejection probability of (5) for $\alpha_{pre} = 1$ (no pretest), $\alpha_{pre} = 0.10$, $\alpha_{pre} = 0.05$ and $\alpha_{pre} = 0.01$ as a function of δ for selected values of ρ , based on 50,000 Monte Carlo draws and 1,000 step random walk approximations to Wiener processes. None of these tests comes close to controlling size uniformly over δ , with null rejection probabilities of at least 20% for some value of (ρ, δ) . Pre-testing for a break seems to both substantially exacerbate and shift the size control problem to larger values of δ .

In fact, to obtain a 5% level test in the no pretest case, one must employ a critical value of approximately 4.9 instead of $\Phi(0.975) \simeq 1.96$ for the t-statistic, with size still approximately equal to the nominal 5% level at the point $(\rho, \delta) = (0.85, 6.8)$. Denote this size corrected test φ_{LS} . Because the worst size distortion occurs for a strictly positive break magnitude, the size corrected hybrid subsampling method of Andrews and Guggenberger (2007a, 2007b) reduces to the size corrected fixed critical value test φ_{LS} . Alternatively, one could always use the test $\varphi_{0.85}(G) = \mathbf{1}[|G(1) - G(0.85)|/\sqrt{0.15} > 1.96]$ which corresponds to a usual 5% level t-test based on the last 15% of the data. As one would expect, though, the power of both φ_{LS} and $\varphi_{0.85}$ is quite poor (we provide power computations in Section 4 below), motivating the construction of an alternative test.

2.1.3 Approximately Weighed Average Power Maximizing Test

No uniformly most powerful test exists for inference about β in absence of knowledge of ρ and δ . Some tests will have good power for certain ranges of value of ρ and δ , while performing poorly for other values. To obtain a reasonable assessment of the overall quality of tests, we therefore




Figure 2: Asymptotic null rejection probabilities of nominal 5% level tests for post break parameter value based on least squares break date estimator.

adopt a weighted average power criterion, as, for instance, in Wald (1943) and Andrews and Ploberger (1994). Specifically, we seek to determine a 5% level test φ that comes close to maximizing

$$\text{WAP}(\varphi) = \int E_{\theta}[\varphi(G)]dF(\theta) \quad (7)$$

where the weighting function F is chosen as follows.

Condition 1 *Under F , ρ is distributed uniform over $[0.15, 0.85]$ and (β, δ) is bivariate normal independent of ρ , with $\beta \sim \mathcal{N}(0, \sigma_{\beta}^2)$, $\delta \sim \mathcal{N}(0, \sigma_{\delta}^2)$ and β independent of $\beta + \delta$, where $\sigma_{\beta}^2 = 22$ and $\sigma_{\delta}^2 = 400$.*

The Gaussianity of the marginal distribution of β and δ in F under Condition 1 is chosen for analytical convenience, and the independence of the pre and post break parameter values β and $\beta + \delta$ mirrors the independence of the pre and post break sample information $G(\rho)$ and $G(1) - G(\rho)$. The value of $\sigma_{\beta}^2 = 22$ is motivated by King's (1988) discussion of overall reasonable point-optimal tests, since it turns out that for $\sigma_{\beta}^2 = 22$, the best 5% level test has approximately 50% weighted average power. The uniform weighting over ρ accords to the choice in Andrews and Ploberger (1994) and is intended to generate reasonable power for all break dates $\rho \in [0.15, 0.85]$. Finally, the value $\sigma_{\delta}^2 = 400$ is motivated as follows: For large δ , (say, $|\delta| > 12$ or so) there is a lot of information about the break date ρ , and good tests can come close to performing as well as if ρ was known. By putting enough weight on rather large values of δ , the choice $\sigma_{\delta}^2 = 400$ ensures that tests that perform well according to WAP share this desirable feature. At the same time, the distribution $\mathcal{N}(0, 400)$ concentrates about half of its mass on $|\delta| < 12$, so tests with high WAP must also perform reasonably in the arguably empirically relevant region for δ where the lack of knowledge of the break date severely complicates good inference about β .

By Girsanov's Theorem, the Radon-Nikodym derivative of the measure of G in (1) with parameter $\theta \in \Theta_0 \cup \Theta_1$ relative to the measure ν of the standard Wiener process W , evaluated at G , is given by

$$f_{\theta}(G) = \exp[\delta G(\rho) + \beta G(1) - \frac{1}{2}(\beta^2 + 2\beta\delta\rho + \delta^2\rho)]. \quad (8)$$

Expressed in terms of the density f_{θ} , the hypothesis test (2) becomes

$$\begin{aligned} H_0 &: \text{The density of } G \text{ is } f_{\theta}, \theta \in \Theta_0 \\ H_1 &: \text{The density of } G \text{ is } f_{\theta}, \theta \in \Theta_1 \end{aligned} \quad (9)$$

and, as noted above, both the null and alternative hypothesis involve the two nuisance parameters δ and ρ . Note, however, that weighted average power can be rewritten as

$$\text{WAP}(\varphi) = \int E_{\theta}[\varphi(G)]dF(\theta) = \int \int \varphi(G)f_{\theta}dF(\theta)d\nu$$

as Fubini’s Theorem justifies the change of order of integration. Thus, maximizing weighted average power is equivalent to maximizing power against the single alternative

$$H_F : \text{The density of } G \text{ is } h = \int f_\theta dF(\theta) \quad (10)$$

and the remaining challenge is to deal with the composite nature of the null hypothesis. The key insight that allows us to make further progress is Lemma 2 of Müller and Watson (2008a),³ which we reproduce here for convenience.

Lemma 1 *Let Λ be a probability distribution on Θ_0 , and let φ_Λ be the best level α test of the null hypothesis H_Λ : The density of G is $\int f_\theta d\Lambda(\theta)$ against H_F . Then for any level α test φ of H_0 against H_F , $\int \varphi_\Lambda h d\nu \geq \int \varphi h d\nu$.*

Proof. Since φ is a level α test of H_0 , $\int \varphi f_\theta d\nu \leq \alpha$ for all $\theta \in \Theta_0$. Therefore, $\int \int \varphi f_\theta d\nu d\Lambda(\theta) = \int \int \varphi f_\theta d\Lambda(\theta) d\nu \leq \alpha$ (where the change in the order of integration is allowed by Fubini’s Theorem), so that φ is also a level α test of H_Λ against H_1 . The result follows by the definition of a best test. ■

Lemma 1 formalizes the perfectly intuitive result that replacing the composite null hypothesis H_0 with the single mixture null hypothesis H_Λ can only simplify the testing problem in the sense of allowing for more powerful tests. Its appeal lies in the fact that the best test of H_Λ against H_F is easily constructed: by the Neyman-Pearson Lemma, the best test rejects for large values of the likelihood ratio statistic $h / \int f_\theta d\Lambda(\theta)$. Thus, Lemma 1 provides a set of explicit power bounds on the original problem, indexed by the distribution Λ .

Lemma 1 can usefully be thought of as generalizing a standard result concerning tests with a composite null hypothesis; see, for instance, Theorem 3.8.1 of Lehmann and Romano (2005): A distribution Λ^{**} is *least favorable* if the best level α test of $H_{\Lambda^{**}}$ against the single alternative H_F is also of level α in the testing problem with the composite null hypothesis H_0 against H_F , so that—using the same reasoning as in the proof of Lemma 1—this test is also the best test of H_0 against H_F . In contrast to this standard result, Lemma 1 is formulated without any restriction on the probability distribution Λ . This is useful because in many contexts, it is difficult to identify the least favorable distribution Λ^{**} (and it may not even exist).

The strategy suggested by Lemma 1 is to work instead with a numerically determined *approximately least favorable* distribution Λ^* : Suppose we knew of a Λ^* such that (i) the best 5% level test of H_{Λ^*} against H_F , φ_{Λ^*} , has power 51.5% and (ii) the best 4.7% level test $\tilde{\varphi}_{\Lambda^*}$ of H_{Λ^*}

³The same insight was applied independently by Andrews, Moreira, and Stock (2007) in the weak instrument problem.

against H_F has power 49.0% and $\tilde{\varphi}_{\Lambda^*}$ is a 5% level test of H_0 . Using Lemma 1, this would imply that no 5% level test of H_0 against H_1 can exist with weighted average power larger than 51.5%, and at the same time, we would have identified the 5% level test $\tilde{\varphi}_{\Lambda^*}$ of H_0 against H_1 whose weighted average power is only 2.5% lower than this bound.

This is the basic idea underlying the construction of our suggested test φ^* detailed in Section 3. For reasons discussed there, the test φ^* is only roughly equivalent to a Neyman-Pearson test $\tilde{\varphi}_{\Lambda^*}$ relative to an approximately least favorable distribution. This is irrelevant for the logic outlined above, only the validity of the power bound via Lemma 1 depends on its derivation from a best (i.e. Neyman-Pearson) test.

We postpone further details of the construction of the power bound and the test φ^* to Section 3 below. The following proposition summarizes the main finding.

Proposition 1 *Under Condition 1,*

(i) *any 5% level test φ of (2) has $WAP(\varphi) \leq \bar{\pi} \simeq 51.5\%$;*

(ii) *the (nonrandomized) test φ^* defined in the appendix is of level $\alpha^* \simeq 5\%$, and has $WAP(\varphi^*) = \pi^* \simeq 49.0\%$.*

The numbers for $\bar{\pi}$, α^* and π^* in Proposition 1 are estimated based on 50,000 independent Monte Carlo draws. As discussed in detail in Section 3, they are *not* subject to any other qualifications, such as a replacement of the level requirement (3) by a finite set of values for (δ, ρ) or a finite step Random Walk approximation for draws of G .

2.2 Multivariate Limiting Problem

Now assume that in the hypothesis testing problem (2) we observe a multivariate version of (1), with

$$\begin{pmatrix} G(s) \\ \tilde{G}(s) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} W(s) \\ \tilde{W}_k(s) \end{pmatrix} + \int_0^s \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{f}(\lambda) \end{pmatrix} d\lambda \quad (11)$$

where \tilde{G} is $k \times 1$, the $k \times 1$ vector A_{21} and the $k \times k$ full rank matrix A_{22} are known, \tilde{W}_k is a k vector standard Wiener process independent of W , and $\tilde{f}(\lambda)$ is a $D_{[0,1]} \mapsto \mathbb{R}^k$ cadlag function, so that $(G, \tilde{G})' \in D_{[0,1]}^{k+1}$. Except for $\text{Var}[G(1)] = 1$, the transformation by A_{21} and A_{22} allows for any full rank covariance matrix of $(G(1), \tilde{G}(1))'$. With $(G, \tilde{G})'$ as the observation, tests φ are now mappings from $D_{[0,1]}^{k+1} \mapsto [0, 1]$, and the rejection probability is given by $E_{\theta, \tilde{f}}[\varphi(G, \tilde{G})]$ (the rejection probability also depends on A_{21} and A_{22} , but we omit this dependence for ease of notation).

For the link to the post break parameter inference problem, think again of a Gaussian location problem

$$\begin{pmatrix} y_t \\ \tilde{y}_t \end{pmatrix} = \begin{pmatrix} \mu + \Delta \mathbf{1}[t \leq \tau] \\ \tilde{\mu}_t \end{pmatrix} + \varepsilon_t, \quad \varepsilon_t \sim \mathcal{N}(0, \Sigma) \quad \text{with } \Sigma = \begin{pmatrix} 1 & A'_{21} \\ A_{21} & A_{21}A'_{21} + A_{22}A'_{22} \end{pmatrix} \quad (12)$$

where $\tilde{y}_t, \tilde{\mu}_t \in \mathbb{R}^k$, so that partial sums $T^{-1/2} \sum_{t=1}^s (y_t, \tilde{y}_t)'$ have the same distribution as $(G, \tilde{G})'$ for $s \in \{t/T\}_{t=1}^T$, where $\rho = \tau/T$, $\beta = T^{1/2}\mu$, $\delta = T^{1/2}\Delta$ and $f(t/T) = T^{1/2}\tilde{\mu}_t$.

The first element in (11), G , is distributed exactly as in the scalar problem (1) above. Thus, if we define φ^* to be the same test as above, that is, φ^* is a function of G only, then φ^* continues to have the properties indicated in Proposition 1. The rejection probability of φ^* obviously does not depend on the distribution of \tilde{G} , so that in particular

$$E_{\theta, \tilde{f}}[\varphi^*(G)] \leq \alpha^* \text{ for all } \tilde{f} \text{ and } \theta \in \Theta_0 \quad (13)$$

$$\inf_{\tilde{f}(\cdot) \in D_{[0,1]}^k} \text{WAP}_{\tilde{f}}(\varphi^*) = \sup_{\tilde{f}(\cdot) \in D_{[0,1]}^k} \text{WAP}_{\tilde{f}}(\varphi^*) = \pi^* \quad (14)$$

where for a generic test $\varphi : D_{[0,1]}^{k+1} \mapsto [0, 1]$, $\text{WAP}_{\tilde{f}}(\varphi) = \int E_{\theta, \tilde{f}}[\varphi(G, \tilde{G})] dF(\theta)$. In other words, φ^* is a robust test of $H_0 : \theta \in \Theta_0$ from observing (11) in the sense that its size and weighted average power does not depend on the nuisance parameter \tilde{f} .

In the following, we restrict attention to the case where \tilde{f} is known to be of the form $\tilde{f}(\lambda) = \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho]$ for $\tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$. This corresponds to a Gaussian location problem (12) where \tilde{y}_t undergoes a single mean shift at the same time as y_t . Write $E_{\theta, \tilde{\beta}, \tilde{\delta}}$ for the expectation with respect to the distribution of $(G, \tilde{G})'$ with \tilde{f} of this form, so that a 5% level test $\varphi : D_{[0,1]}^{k+1} \mapsto [0, 1]$ now satisfies $E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi(G, \tilde{G})] \leq 0.05$ for all $\theta \in \Theta_0$, $\tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$.⁴

Proposition 2 (i) For any unbiased 5% level test φ ,

$$E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_\rho(G)] - E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi(G, \tilde{G})] \geq 0 \text{ for all } \theta \in \Theta_1, \tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$$

where $\varphi_\rho(G) = \mathbf{1}[|\frac{G(1)-G(\rho)}{\sqrt{1-\rho}}| > \Phi(0.975)]$.

(ii) Let \tilde{F} be the probability distribution for $(\theta, \tilde{\beta}, \tilde{\delta})$ with the same marginal distribution for θ as under Condition 1, and $(\tilde{\beta}, \tilde{\delta}) = (\beta A_{21}, \delta A_{21})$ conditional on θ . For any 5% level test

$$\int E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi(G, \tilde{G})] d\tilde{F}(\theta, \tilde{\beta}, \tilde{\delta}) \leq \bar{\pi} \simeq 51.5\%.$$

⁴All following propositions also hold for the scalar case ($k = 0$) with the natural interpretation of notation.

(iii) Suppose in addition that $\tilde{\delta} = 0$ known, so that tests $\varphi : D_{[0,1]}^{k+1} \mapsto [0, 1]$ are of 5% level if $E_{\theta, \tilde{\beta}, 0}[\varphi(G, \tilde{G})] \leq 0.05$ for all $\theta \in \Theta_0$ and $\tilde{\beta} \in \mathbb{R}^k$. Consider the group of transformations

$$\begin{pmatrix} G(s) \\ \tilde{G}(s) \end{pmatrix} \rightarrow \begin{pmatrix} G(s) \\ \tilde{G}(s) + \tilde{b}s \end{pmatrix}, \tilde{b} \in \mathbb{R}^k. \quad (15)$$

For any 5% level test φ invariant to (15), the rejection probability $E_{\theta, \tilde{\beta}, 0}[\varphi(G, \tilde{G})]$ does not depend on $\tilde{\beta}$, and under Condition 1,

$$WAP_{inv}(\varphi) = \int E_{\theta, \tilde{\beta}, 0}[\varphi(G, \tilde{G})] dF(\theta) \leq \bar{\pi}_{inv}(\bar{R}^2)$$

where $\bar{R}^2 \geq R^2 = A'_{21}(A_{21}A'_{21} + A_{22}A'_{22})^{-1}A_{21}$, and $\bar{\pi}_{inv}(0) = \bar{\pi} \simeq xx$, $\bar{\pi}_{inv}(0.2) \simeq xx$, $\bar{\pi}_{inv}(0.4) \simeq xx$, $\bar{\pi}_{inv}(0.6) \simeq xx$, and $\bar{\pi}_{inv}(0.8) \simeq xx$.

Part (i) of Proposition 2 establishes that the usual two-sided t-test using post break data $G(1) - G(\rho)$, φ_ρ , is the uniformly most powerful unbiased test in the multiparameter problem with (11) observed. With ρ unknown this test is infeasible. It nevertheless provides an arguably relevant benchmark to assess the performance of other tests. In particular, note that φ_ρ , just like φ^* , does not depend on \tilde{G} , but is a function of G only. We compare the power of φ^* to that of φ_ρ in detail in Section 4 below, and find that for large $|\delta|$, the power becomes quite close. Thus, for large $|\delta|$, φ^* has similar rejection properties as the uniformly most powerful unbiased test, so that there is at most a small cost in terms of power for ignoring the additional information in \tilde{G} . (Although φ^* is not unbiased, as it is not similar—see Section 3 below.)

Part (ii) of Proposition 2 shows that for a particular weighting function \tilde{F} with the same marginal distribution as F on θ , no 5% level test can have weighted average power larger than $\bar{\pi}$. At the same time, (14) implies that the weighted average power of φ^* with respect to \tilde{F} is equal to π^* , and, by Proposition 1, π^* is only slightly smaller than $\bar{\pi}$. In this sense, φ^* is close to being admissible also in the hypothesis testing problem with (11) observed. Thus, any test with "substantially" larger power than φ^* for some $(\theta, \tilde{\beta}, \tilde{\delta})$ is less powerful for some other $(\theta, \tilde{\beta}, \tilde{\delta})$. Given this inevitable trade-off, the choice of φ^* of approximately maximizing weighted average power relative to \tilde{F} has the perhaps appealing implication of yielding a test whose weighted average power over θ does not depend on the nuisance parameters $(\tilde{\beta}, \tilde{\delta})$.

Finally, part (iii) of Proposition 2 provides bounds on the gains in weighted average power that are possible if it is known that the other parameters do not undergo a break, $\tilde{\delta} = 0$ (or, equivalently, if $\tilde{\delta}$ was known) under an invariance requirement that corresponds to changes of the mean of the additional k observations. A maximal invariant to this group is given by $(G(s), \tilde{G}(s) - s\tilde{G}(1))$. Intuitively, these power gains are possible because if \tilde{G} is correlated with

$G (R^2 > 0)$, observing $\tilde{G}(s) - s\tilde{G}(1) = \tilde{W}(s) - s\tilde{W}(1)$ provides information about $W(s) - sW(1)$, which is useful for learning about ρ and δ . In the extreme case of perfect correlation ($R^2 = 1$), $W(s) - sW(1)$ is effectively observed, and by comparing $G(s) - sG(1)$ to $W(s) - sW(1)$, one can back out ρ and δ and obtain as powerful inference about β as if there was no break in the model. The numbers for $\bar{\pi}_{inv}$ were estimated using xx,000 independent Monte Carlo draws. They show that rather large values of R^2 are necessary before knowledge of $\tilde{\delta} = 0$ could potentially be used to generate tests with substantially larger weighted average power than φ^* .

In summary, the test φ^* that disregards the additional observation \tilde{G} in (11) is (i) robust in the sense of providing reliable inference about β regardless of the nuisance function \tilde{f} ; (ii) for large δ has a rejection profile that is close to that of the uniformly most powerful unbiased test; (iii) is close to admissible in a model where other parameters undergo a break at the same time; (iv) approximately maximizes weighted average power even when it is known that only the parameter of interest is subject to a break as long as R^2 is small. For these reasons, the remainder of this paper focusses on φ^* as a reasonable test also in the multiparameter context of (11). We note, however, that it should be possible in principle to use the numerical methods discussed in Section 3 to also identify 5% level feasible test that comes close to maximizing weighted average power by efficiently exploiting the additional observation \tilde{G} when it is known that only the parameter of interest is subject to a break, for any particular value of R^2 .

2.3 Asymptotic Efficiency Implications for Underlying Models

In this subsection, we discuss how the scalar and multiparameter limiting problems (1) and (11) arise as the relevant asymptotic problems in standard small sample post break parameter inference. We consider two distinct approaches to asymptotic efficiency of tests: On the one hand, we rely on LeCam's Limits of Experiments theory to derive upper bounds on power in correctly specified likelihood models. On the other hand, we exploit the recent results of Müller (2007) and provide upper bounds on post break parameter tests in time series GMM models under an asymptotic robustness constraint.

2.3.1 Parametric Models

Let $X_T = (x_{T,1}, \dots, x_{T,T}) \in \mathbb{R}^{qT}$ be the available data in a sample of size T , so that tests φ_T are sequences of (measurable) functions $\varphi_T : \mathbb{R}^{qT} \mapsto [0, 1]$. The density of X_T , relative to some σ -finite measure μ_T , is given by the product $\prod_{t=1}^T f_{T,t}(\Gamma_t)$, when the parameter $\Gamma \in \mathbb{R}^{k+1}$ takes on the value Γ_t at time t . This form of the likelihood arises naturally when $f_{T,t}(\Gamma_t)$ is the density of $x_{T,t}$ conditional on $\mathfrak{F}_{T,t-1}$, the σ -field generated by $\{x_{T,s}\}_{s=1}^{t-1}$. We will refer to the model with

density $\prod_{t=1}^T f_{T,t}(\Gamma_0)$ as the 'stable' model. Define $l_{T,t}(\Gamma) = \ln f_{T,t}(\Gamma)$, $s_{T,t}(\Gamma) = \partial l_{T,t}(\Gamma)/\partial \Gamma$ and $h_{T,t} = \partial s_{T,t}(\Gamma)/\partial \Gamma'$, and suppose $T^{-1} \sum h_{T,t} \xrightarrow{p} -H$ in the stable model for some positive definite and known $(k+1) \times (k+1)$ matrix H , so that H^{-1} is the Fisher information. In the unstable model, the parameter vector Γ evolves as

$$\Gamma_{T,t} = \Gamma_0 + T^{-1/2} \begin{pmatrix} \beta \omega \\ \tilde{\beta} \end{pmatrix} + T^{-1/2} \begin{pmatrix} \delta \omega \\ \tilde{\delta} \end{pmatrix} \mathbf{1}[t \leq \rho T] \quad (16)$$

where ω^2 is the 1-1 element of H , $\beta, \delta, \rho \in \mathbb{R}$ and $\tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$, and $\rho \in [0.15; 0.85]$. We assume Γ_0 to be known; while this unrealistic, such knowledge can only increase the upper bounds on power derived in Proposition 3 below. The hypothesis of interest continuous to concern the value of the post break parameter

$$H_0 : \beta = 0 \quad \text{against} \quad H_1 : \beta \neq 0. \quad (17)$$

The factor $\omega > 0$ in (16) ensures that asymptotically, the small sample problem (17) maps to the limiting problem (11) where the disturbance in G is a *standard* Wiener process. Tests $\varphi_T : \mathbb{R}^{qT} \mapsto [0, 1]$ of (17) have unconditional rejection probability $E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_T(X_T)]$ in a sample of size T , and φ_T is defined to be of asymptotic level 5% if $\limsup_{T \rightarrow \infty} E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_T(X_T)] \leq 5\%$ for all $\theta \in \Theta_0$, $\tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$.

Under suitable regularity conditions on the parametric model, one can show that the likelihood ratio statistic LR_T between the model with parameter evolution described by (16) and the stable model satisfies

$$\begin{aligned} \ln \text{LR}_T &= T^{-1/2} \sum_{t=1}^T (\Gamma_{T,t} - \Gamma_0)' s(\Gamma_0) + \frac{1}{2} T^{-1} \sum_{t=1}^T (\Gamma_{T,t} - \Gamma_0)' h_t(\Gamma_0) (\Gamma_{T,t} - \Gamma_0) + o_p(1) \\ &\Rightarrow \int_0^1 \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix}' \Sigma^{-1} d \begin{pmatrix} G(\lambda) \\ \tilde{G}(\lambda) \end{pmatrix} - \frac{1}{2} \int_0^1 \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix}' \Sigma^{-1} \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix} d\lambda \end{aligned} \quad (18)$$

where $\Sigma = \omega^2 H^{-1}$. The r.h.s. of (18) may be recognized as log of the Radon-Nikodym derivative of the distribution of $(G, \tilde{G})'$ with respect to the distribution of the $(k+1) \times 1$ Wiener process $(W, (A_{21}W + A_{22}\tilde{W}_k)')$. This suggests that the information regarding $(\theta, \tilde{\beta}, \tilde{\delta})$ from observing X_T converges in large samples to that contained in $(G, \tilde{G})'$. This intuition is made formally precise in the Limit of Experiments theory pioneered by LeCam; see, for instance, van der Vaart (1998) for an introduction.

Condition 2 in the appendix states sufficient regularity assumptions that imply convergence of experiments in this sense. The Asymptotic Representation Theorem (see, for instance, Theorem 9.3 of van der Vaart (1998)) and Prohorov's Theorem then ensure that for any test

$\varphi_T : \mathbb{R}^{qT} \mapsto [0, 1]$ and any subsequence T' of T , there exists a further subsequence T'' of T' and a test $\varphi : D_{[0,1]}^{k+1} \mapsto [0, 1]$ such that

$$E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_{T''}(X_{T''})] \rightarrow E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi(G, \tilde{G})] \text{ for all } \theta \in \Theta_0 \cup \Theta_1, \tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k. \quad (19)$$

Thus, along any such subsequence T'' , the asymptotic properties of φ_T are exactly equivalent to a particular test φ in the limiting problem discussed in Section 2.2 above. Define a test φ_T as *asymptotically unbiased* if any test φ satisfying (19) is unbiased. Similarly, define a test φ_T as *asymptotically invariant* if any test φ satisfying (19) is invariant to (15).

Proposition 3 (i) *For any asymptotically unbiased test φ_T of asymptotic level 5%,*

$$\limsup_{T \rightarrow \infty} E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_T(X_T)] \leq E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_\rho(G)] \text{ for all } \theta \in \Theta_1, \tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$$

(ii) *For any test φ_T of asymptotic level 5%,*

$$\limsup_{T \rightarrow \infty} \int E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_T(X_T)] d\tilde{F}(\theta) \leq \bar{\pi}.$$

(iii) *For any asymptotically invariant test φ_T of asymptotic level 5%,*

$$\limsup_{T \rightarrow \infty} \int E_{\theta, \tilde{\beta}, 0}[\varphi_T(X_T)] dF(\theta) \leq \pi_{inv}(\bar{R}^2) \text{ for all } \tilde{\beta} \in \mathbb{R}^k.$$

Proposition 3 almost exactly mirrors the results of Proposition 2 above. Furthermore, we argue in Section 4 below that the feasible test $\hat{\varphi}^* : \mathbb{R}^{qT} \mapsto [0, 1]$ has the same asymptotic rejection properties as the test φ^* under (16). Thus, to the extent that Proposition 2 implies φ^* to be an attractive test in the context of the multivariate limiting problem (11), $\hat{\varphi}^*$ is correspondingly an attractive test in the context of post break inference in a correctly specified parametric model that is subject to a local break.

2.3.2 GMM Models

We now turn to an application of the asymptotic efficiency concept introduced by Müller (2007), which we use here to state asymptotic power bounds on post break parameter inference in a class of GMM models. Let the moment condition be the \mathbb{R}^l -valued function $g(\cdot, \cdot)$, so that $E[g(x_{T,t}, \Gamma_0)] = 0$ when the true parameter at date t is given by $\Gamma = \Gamma_0$. Let the $\mathbb{R}^{(k+1) \times l}$ valued function $\Upsilon(\cdot, \Gamma)$ be the partial derivative of g with respect to Γ , and write $g_t(\Gamma) = g(x_{T,t}, \Gamma)$ and $\Upsilon_t(\Gamma) = \Upsilon(x_{T,t}, \Gamma)$ for notational simplicity. Under the parameter evolution (16), conditional on

τ , the natural estimators for the pre and post break value of Γ are given by the GMM estimator using pre and post break data,

$$\begin{aligned}\hat{\Gamma}_{pre}(\tau) &= \arg \min_{\Gamma} \left(\sum_{t=1}^{\tau} g_t(\Gamma) \right)' \hat{V}_{pre} \left(\sum_{t=1}^{\tau} g_t(\Gamma) \right) \\ \hat{\Gamma}_{post}(\tau) &= \arg \min_{\Gamma} \left(\sum_{t=\tau+1}^T g_t(\Gamma) \right)' \hat{V}_{post} \left(\sum_{t=\tau+1}^T g_t(\Gamma) \right)\end{aligned}$$

with \hat{V}_{pre} and \hat{V}_{post} possibly data dependent $l \times l$ positive definite weighting matrices, and associated covariance matrix estimators $\hat{\Omega}_{pre}(\tau)$ and $\hat{\Omega}_{post}(\tau)$ of $\hat{\Gamma}_{pre}(\tau)$ and $\hat{\Gamma}_{post}(\tau)$, respectively. A natural starting point for inference about the post break parameter value in absence of knowledge of $\rho \in [0.15; 0.85]$ are thus the processes $\{\hat{\Gamma}_{pre}(\lfloor \cdot T \rfloor), \hat{\Gamma}_{post}(\lfloor \cdot T \rfloor), \hat{\Omega}_{pre}(\lfloor \cdot T \rfloor), \hat{\Omega}_{post}(\lfloor \cdot T \rfloor)\} \in D_{[0.15, 0.85]}^{4(k+1)}$, where $\lfloor x \rfloor$ denotes the largest integer smaller or equal to $x \in \mathbb{R}$. Assuming that $T^{-1/2} \sum_{t=1}^{\lfloor \cdot T \rfloor} g_t(\Gamma_0) \Rightarrow \Sigma_g^{1/2} W_{k+1}(\cdot)$ for some positive definite matrix Σ_g and $T^{-1} \sum_{t=1}^{\lfloor \cdot T \rfloor} \Upsilon_t(\Gamma_0) \xrightarrow{p} \bar{\Upsilon}$ for some full column rank matrix $\bar{\Upsilon}$, the usual Taylor expansion arguments yield, for the case of efficient GMM estimation,

$$\begin{aligned}\{T^{1/2} \hat{\Gamma}_{pre}(\lfloor sT \rfloor), T^{1/2} \hat{\Gamma}_{post}(\lfloor sT \rfloor), T \hat{\Omega}_{pre}(\lfloor sT \rfloor), T \hat{\Omega}_{post}(\lfloor sT \rfloor)\} \\ \Rightarrow \left\{ \begin{pmatrix} G(s) \\ \tilde{G}(s) \end{pmatrix}, \begin{pmatrix} G(1) - G(s) \\ \tilde{G}(1) - \tilde{G}(s) \end{pmatrix}, s\omega^2\Sigma, (1-s)\omega^2\Sigma \right\}\end{aligned}\quad (20)$$

in $D_{[0.15, 0.85]}^{(k+1) \times 4}$, where now $\omega^2\Sigma = (\bar{\Upsilon}'\Omega_g^{-1}\bar{\Upsilon})^{-1}$ (cf. Andrews (1993)).

Consider the set of data generating processes for X_T that satisfy (20). One might want to choose the tests φ_T of (17) in a way that whenever (20) holds with $\beta = 0$, the test does not overreject asymptotically, that is $\limsup_{T \rightarrow \infty} E_{\theta, \tilde{\beta}, \tilde{\delta}}[\varphi_T(X_T)] \leq 0.05$ for all $\theta \in \Theta_0$, $\tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$ for *any* sequence of distributions of X_T that satisfies (20). Müller (2007) shows that under this robustness constraint, the best small sample test becomes the best test in the limiting problem (that is, with the r.h.s. of (20) assumed observed), evaluated at sample analogues (that is, at $\{T^{1/2} \hat{\Gamma}_{pre}(\lfloor \cdot T \rfloor), T^{1/2} \hat{\Gamma}_{post}(\lfloor \cdot T \rfloor), T \hat{\Omega}_{pre}(\lfloor \cdot T \rfloor), T \hat{\Omega}_{post}(\lfloor \cdot T \rfloor)\}$). Proposition 2 above shows in which sense φ^* may be considered approximately best in the equivalent problem of observing $(G(\cdot), \tilde{G}(\cdot))' \in D_{[0,1]}^{k+1}$ directly. Thus, part (ii) of Proposition 2 implies, for instance, that the weighted average asymptotic power (defined as in part (ii) of Proposition 3) of any 5% level test is no larger than $\bar{\pi}$. We discuss in detail in Section 4 below how to construct the small sample analogue $\hat{\varphi}^*$, and show in Proposition 4 that its asymptotic rejection profile is equal to the rejection profile of φ^* whenever (20) holds, so its asymptotic weighted average power is equal to π^* . Thus, $\hat{\varphi}^*$ is the approximately best test among all robust tests in the sense of Müller (2007). We omit a formal statement analogous Proposition 3 to conserve space.

3 Determination of an Approximately Efficient Test with Uniform Size Control

This section describes the methods and algorithm that underlie the claim of Proposition 1, that is we discuss the determination of a 5% level test of (2) based on the observation G as in (1) that approximately maximizes weighted average power with a weighting function as described in Condition 1.

As outlined in subsection 2.1.2 above, the key challenge is to identify an approximately least favorable distribution over the two dimensional nuisance parameter of the break date and break magnitude. The first subsection provides details on the numerical algorithm for obtaining this distribution, which is a suitably modified version of what is developed in Müller and Watson (2008a) for the problem considered here. The second subsection discusses the relationship and properties of the least favorable distribution from a decision theoretic and Bayesian perspective. In the third subsection, we introduce a new approach to the numerical study of size control of tests of a composite null hypothesis, which underlies our claim of uniform size control of the test φ^* . Finally, the fourth subsection contains details on the Monte Carlo determination of the power bound, which is complicated by the impossibility of generating and representing (pseudo) random Wiener process draws on a discrete computer.

3.1 Approximately Least Favorable Distribution

The guiding principle for the construction of an appropriate approximately least favorable distribution Λ^* is following property discussed in subsection 2.1.2: for a distribution Λ to be approximately least favorable, it must be true that a slight adjustment of the critical value (or, equivalently, of the level of the best test of H_Λ against H_F) yields a 5% level test under H_0 . A violation of this property guides how Λ should be amended, with more mass on those values of θ where the overrejection occurred. By suitably iterating changes of Λ with computations of its rejection profile under H_0 , an approximately least favorable Λ^* can be determined numerically.

Müller and Watson (2008a) directly implement this idea by considering distributions Λ that have mass on a finite number of points $\{\theta_1, \dots, \theta_L\} \subset \Theta_0$. If for a given Λ , φ_Λ overrejects at $\theta' \in \Theta_0$, then $\theta_{L+1} = \theta'$ is added to the list. A new Λ is determined by assigning the appropriate probabilities on $\{\theta_1, \dots, \theta_{L+1}\}$ to ensure exact size control at $\{\theta_1, \dots, \theta_{L+1}\}$, and the algorithm iterates these steps until an appropriate least favorable distribution is found. The numerical feasibility of this procedure depends on L not to become too large (say, $L < 40$ or so).

For the problem studied here, this strategy turns out to not work well: If φ_Λ overrejects for

a moderately large value of $|\delta|$ and a (possibly small) interval of values for ρ , then the inclusion of any single point $\theta' = (0, \delta', \rho') \in \Theta_0$ in Λ does not remedy the overrejection problem for all values of ρ in the interval. The reason is that for moderately large values of $|\delta|$, the distribution of G with ρ'' slightly different from ρ' is already too dissimilar for size control at ρ' to imply approximately size control at ρ'' . Roughly speaking, with large breaks, different break dates lead to almost singular probability distributions for G , so that a very large number of points would be required to ensure approximate overall size control.

Instead, we note that for Λ to be approximately least favorable, an equivalent condition is that (the slightly level adjusted) test φ_Λ is of 5% level under any H_Ψ : The density of G is $\int f_\theta d\Psi(\theta)$, for all probability distributions Ψ on Θ_0 . By initially restricting this condition to a finite set $\mathcal{P} = \{\Psi_1, \dots, \Psi_M\}$ with moderately large M , the above algorithm can be successfully implemented to identify Λ_M^* , the least favorable mixture of \mathcal{P} , such that $\varphi_{\Lambda_M^*}$ controls size for all H_Ψ , $\Psi \in \mathcal{P}$. Collect the distributions Ψ_j that receive positive mass by Λ_M^* in $\mathcal{P}^* = \{\Psi_1^*, \dots, \Psi_M^*\} \subset \mathcal{P}$. Using Λ_M^* as the new starting point, consider a further set $\mathcal{P}' = \{\Psi'_1, \dots, \Psi'_{M'}\}$ and again identify $\Lambda_{M'}^*$, a mixture of \mathcal{P}^* and \mathcal{P}' , so that $\varphi_{\Lambda_{M'}^*}$ controls size for all H_Ψ , $\Psi \in \mathcal{P}^* \cup \mathcal{P}'$. This procedure is then iterated, and by considering sets of probability distributions that more and more closely approximate point masses in Θ_0 (although they still average over short intervals of ρ), a suitable Λ^* can be determined. In our implementation, we choose Ψ 's with (not necessarily mean zero) Gaussian distributions over δ conditional on ρ , with smaller and smaller variances in later steps of the iteration.

We relegate details on this algorithm to the appendix, but discuss here two additional issues that are relevant to the determination of the approximate least favorable distribution Λ^* .

On the one hand, the observation G is a transformation of a Wiener process, so that computation of the Neyman-Pearson tests and their rejection profile requires the simulation of (pseudo) random continuous time processes. Discrete time Gaussian random walk approximation based on, say, 1000 steps are computationally expensive and of uncertain accuracy. For this reason, we approximate all integrals over ρ by sums with $\rho \in \{1/100, 2/100, \dots, 1\}$. The rejection probability of the (approximate) Neyman-Pearson test thus becomes a function of

$$\left\{G\left(\frac{l}{100}\right)\right\}_{l=1}^{100}. \quad (21)$$

The 100 random variables (21) can be simulated without any approximation error, as they are exactly jointly Gaussian. What is more, the fact that φ^* only depends on G through (21) significantly simplifies the determination of the size of φ^* discussed in the following subsection.

A power bound on tests that are functions of (21) is not, however, in general a valid power bound on the original problem involving the observation G . The correct implementation of the

power bound via Lemma 1 requires evaluation of the actual density (8) of G , a complication we address in Section 3.4 below. In the end, the comparison of this power bound with the power of the feasible test φ^* reveals whether working with (21) instead of the whole process G might have led to substantial inaccuracies in the resulting approximately least favorable distribution, and Proposition 1 shows that these are small.

On the other hand, we chose to construct the feasible test φ^* by combining two different tests, with a supF-type statistic based on the 100 observations (21) determining to which one it switches: For large values of supF, that is strong evidence for a break, φ^* is essentially identical to the usual two-sided t-test on least-squares estimated post break data, \hat{t} in (6), whereas for small values of supF, φ^* is equal to the Neyman-Pearson test of H_{Λ^*} against H_F . The motivation for this switching rule is threefold: First, it formally justifies the "rule of thumb" that for highly significant realizations of a structural break test statistic, least-squares based inference is just fine, with a 'patch' necessary only for moderate break magnitudes δ . Second, it ensures that even for very large breaks ($|\delta| \rightarrow \infty$), the test has attractive properties—this cannot be ensured by construction by any weighted average power criterion with integrable weighting function, as any such weighting concentrates almost all of its mass on a compact set. Third, the switch to an analytically easily tractable test statistic facilitates the study of the size properties of φ^* for the (unbounded) set of large δ discussed in the next subsection.

The adoption of this switching requires a determination of Λ^* such that the overall test φ^* is of level 5%, rather than the critical value adjusted Neyman-Pearson test $\tilde{\varphi}_{\Lambda^*}$ of H_{Λ^*} against H_F in isolation. While it is true that for large $|\delta|$, the t-test and $\tilde{\varphi}_{\Lambda^*}$ behave quite similarly (after all, for $|\delta|$ large, there is ample information about ρ , so that both $\tilde{\varphi}_{\Lambda^*}$ and the t-test are approximately equal to the best infeasible test with ρ known), we choose a cut-off value for supF that is just large enough to ensure that the t-test controls size when the probability of switching is close to one. Thus, for values of $|\delta|$ where the switch to the t-test only occurs, say, half the time, the t-test has null rejection probability larger than 5%. Given the switching rule, Λ^* must therefore induce a compensating underrejection in this region of δ . In the above algorithm, this is achieved by requiring that the rejection probability of φ_{Λ^*} is substantially lower than 5% under H_{Ψ} for Ψ 's that concentrate on these regions (which in turn leads to a Λ^* with substantial mass on these Ψ 's).

3.2 Decision Theoretic and Bayesian Interpretation

From a decision theoretic perspective, the least favorable prior has a minimax interpretation in the problem of distinguishing between H_0 against H_F . Suppose a false rejection of H_0 induces




Figure 3: Approximate Least Favorable Distribution Λ^*

loss 1, a false acceptance of H_F induces loss $L_F > 0$, and a correct decision has loss 0. Then risk is given by

$$R(\theta, \varphi) = \mathbf{1}[\theta \in \Theta_0]E_\theta[\varphi(G)] + L_F \mathbf{1}[\theta \in \Theta_1] \int E_t[1 - \varphi(G)]dF(t). \quad (22)$$

The level α test based on the least favorable prior Λ^{**} (supposing it exists) minimizes $\sup_{\theta \in \Theta} R(\theta, \varphi)$ among all tests φ for $L_F = \alpha/(1 - WAP(\varphi_{\Lambda^{**}}))$, and achieves $\sup_{\theta \in \Theta} R(\theta, \varphi) = \alpha$. The α^* -level test based on the *approximately* least favorable prior Λ^* achieves $\sup_{\theta \in \Theta} R(\theta, \varphi^*) = \alpha^*$ for $L_F = \alpha^*/(1 - \pi^*)$, and $\inf_\varphi \sup_{\theta \in \Theta} R(\theta, \varphi) \geq \alpha^*(1 - \bar{\pi})/(1 - \pi^*) \simeq 0.951\alpha^*$ by Proposition 1, so φ^* is an approximately minimax decision rule. In this reasoning, the usual requirement of size control becomes an endogenous solution to the desire to find a maximin decision rule under risk (22).

In assessing the appeal of (inherently pessimistic) minimax rules, it is instructive to consider the reasonableness of the implied least favorable distribution. Figure 3 plots the approximately least favorable distribution Λ^* determined by the above algorithm for $\delta \geq 0$; by construction, the distribution is symmetric in the sign of δ . Overall, apart from some peculiarities close to the endpoint for the break date at $\rho = 0.85$, Λ^* does not seem particularly bizarre.

From a Bayesian perspective, one might want to decide between H_0 and H_1 by computing posterior odds. With a prior of F on Θ_1 , and a prior of Λ^* on Θ_0 , φ_{Λ^*} rejects for large values

of the resulting posterior odds, or, equivalently, for large values of the Bayes factor. Apart from the switching to a t-test discussed above, which in this context may be regarded as an approximation to the posterior, an analysis based on φ^* thus also has a straightforward Bayesian interpretation. The prior Λ^* on H_0 and the cut-off value $\simeq 2.41$ are endogenously determined so that Bayes factors above the cut-off value—that is, Bayes factors that occur for at most 5% of draws under H_0 for *all* values of $\theta \in \Theta_0$ —are as frequent as possible under H_F . In this way, the rule of rejecting for a large Bayes factor also has attractive frequentist properties.

It might be instructive to contrast this to the properties of the test that rejects for a large Bayes factor when the prior on the null hypothesis is not endogenized in this fashion. For instance, in this problem, a natural choice for the prior distribution Λ on Θ_0 is a uniform distribution for ρ on $[0.15, 0.85]$, and $\delta \sim \mathcal{N}(0, 400)$ independent of ρ ; this mirrors the prior F of Condition 1 on the alternative Θ_1 , except that $\sigma_\beta^2 = 0$. With that prior, one must choose a cut-off value of $\simeq 10$, rather than 2.41, to ensure that only 5% of the time, one observes Bayes factors larger than the cut-off for *all* values of $\theta \in \Theta_0$. Correspondingly, the probability of rejection under H_F falls from $\pi^* \simeq 49.0\%$ to 36.7% with this choice of Λ and cut-off value.

3.3 Size Control

With a candidate test φ^* determined as outlined in the last subsection, the question is whether it actually controls size, i.e. whether

$$\sup_{\theta \in \Theta_0} E_\theta[\varphi^*(G)] \leq 0.05. \quad (23)$$

By construction of φ^* , this is "approximately" true, but we now discuss how to establish (23) more rigorously, that is part (ii) of Proposition 1.

The problem we face is a standard one: What is the size of a given test under a composite null hypothesis? For some nonstandard hypothesis testing problems, the form of the test greatly simplifies this issue. For instance, the conditional likelihood ratio statistic of Moreira and Andrews, Moreira, and Stock (2006), or the statistic suggested in Jansson and Moreira (2006), have by construction an (asymptotic) null rejection probability that does not depend on the nuisance parameter. For other tests or testing problems, however, essentially nothing analytical is known about the rejection probability. The usual approach then is to resort to a Monte Carlo grid search: Choose a finite set $\Theta_{grid} \subset \Theta_0$, estimate the rejection probability by Monte Carlo for each $\theta \in \Theta_{grid}$, and conclude that the test controls size if the largest of these rejection probabilities is smaller or equal to the nominal level. Examples for this approach include Stock and Watson (1996), Bunzel and Vogelsang (2005), Srikanthakumar and King (2006), Andrews,

Moreira, and Stock (2007), Andrews and Guggenberger (2007a, 2007b, 2007c), Guggenberger (2008), and Müller and Watson (2008a), among others.⁵ For the specific problem here, this approach would amount to computing the Monte Carlo estimate of the null rejection probability

$$\hat{p}(\theta) = N^{-1} \sum_{i=1}^N \varphi^*(G_i) \quad \text{with } G_i(s) = W_i(s) + \delta \min(\rho, s) \quad (24)$$

for all $\theta = (0, \delta, \rho) \in \Theta_{grid}$, where W_i are N (pseudo) random draws of a standard Wiener process⁶ (which may or may not be the same across different θ), and to conclude that φ^* is of level 5% if $\sup_{\theta \in \Theta_{grid}} \hat{p}(\theta) < 0.05$.

Clearly, though, this approach is not fully satisfying. Without some knowledge about the smoothness of $\hat{p}(\theta)$, even a fine grid search of this form does not provide any upper bound on $\sup_{\theta \in \Theta_0} \hat{p}(\theta)$, simply because the rejection probability $\hat{p}(\theta)$ could be very different between the grid points. What is more, Θ_0 is typically unbounded, so that it is impossible from the start to finely cover Θ_0 . We now describe an approach that, while still based on Monte Carlo estimation,⁷ can handle both these difficulties by bounding the function $\hat{p}(\theta)$ on the whole set Θ_0 .

The key idea is the following change of perspective: Rather than to only regard the average $\hat{p}(\theta)$ in (24) as a function of θ , consider instead $\varphi^*(G_i)$ as a function of $\theta = (0, \delta, \rho) \in \Theta_0$, conditional on W_i . The actual data G_i is a function of the (pseudo) randomness W_i and the nuisance parameter θ , so that for each realization of W_i , the test statistic $\varphi^*(G_i)$ may be regarded as a function of θ , $\varphi^*(G_i) = S_i(\theta)$ with $S_i : \Theta_0 \mapsto \{0, 1\}$ (as long as φ^* is non-randomized). The function $S_i(\theta)$ is, of course, known, and one can analytically study its properties. So suppose initially that it was possible to exactly determine and store the "critical regions" $C_i = \{\theta : S_i(\theta) = 1\}$, $i = 1, \dots, N$, that is the range of values for θ for which the test $\varphi^*(G_i)$ rejects.⁸ For instance, if the nuisance parameter space was one dimensional, C_i would typically be a finite union of intervals, and it would suffice to determine their endpoints (possibly including $\pm\infty$). Clearly, the overall Monte Carlo estimate of the null rejection probability function $\hat{p} : \Theta_0 \mapsto [0, 1]$

⁵Dufour (2006) suggests searching for the maximum Monte Carlo rejection probability directly using simulated annealing techniques, which may also fail to yield the global maximum.

⁶As noted above, suffices to generate $\{W(l/100)\}_{l=1}^{100}$, which is actually feasible.

⁷At least in principle, one could rerun this Monte Carlo size algorithm with new random draws before each application of φ^* , so that the Monte Carlo error becomes part of the randomized nature of the actual test. It should be possible to appropriately combine the arguments in Dufour (2006) with what is suggested here to thereby obtain an overall randomized test of *exact* level 5%.

⁸This terminology is meant to be suggestive only; the actual critical region, of course, is the subset of the sample space $D_{[0,1]}$ for which φ^* rejects.

is simply the average of these critical regions

$$\hat{p}(\theta) = N^{-1} \sum_{i=1}^N \mathbf{1}[\theta \in C_i]$$

and size control amounts to $\sup_{\theta \in \Theta_0} \hat{p}(\theta) \leq 0.05$.

Now for our problem, the nuisance parameter space is two dimensional, and the form of φ^* does not make it feasible to exactly describe the sets C_i analytically. So consider instead a finite partition $\mathcal{Q} = \{Q_1, Q_2, \dots, Q_K\}$ of Θ_0 , such as the *rectangles* of a grid, and suppose study of S_i allows one to conclude that the sets $\{Q_j : j \in J_i\} \subset \mathcal{Q}$ with $J_i \subset \{1, 2, \dots, K\}$ cover C_i , i.e. $C_i \subset \bigcup_{j \in J_i} Q_j$. It then suffices to keep track of the index sets J_i , $i = 1, \dots, N$ to establish the bound

$$\sup_{\theta \in \Theta_0} \hat{p}(\theta) \leq \bar{p} = \sup_{\theta \in \Theta_0} N^{-1} \sum_{i=1}^N \mathbf{1}[\theta \in \bigcup_{j \in J_i} Q_j] = \max_j N^{-1} \sum_{i=1}^N \mathbf{1}[j \in J_i] \quad (25)$$

and one can conclude that φ^* is a 5% level test if $\bar{p} \leq 0.05$. Note that this bound remains valid (although it becomes more conservative) when the sets J_i are larger than necessary, that is if $j \in J_i$ despite $C_i \cap Q_j = \emptyset$.

This approach benefits from a finer partition \mathcal{Q} of Θ_0 with K large in two ways. On the one hand, a fine \mathcal{Q} allows for a relatively accurate description of the actual critical region C_i , thus making the bound (25) less conservative. On the other hand, it will typically become easier to decide whether or not $C_i \cap Q_j = \emptyset$, for each $j = 1, \dots, K$. But the computational cost of making K large is also substantial, as the brute force implementation requires a total of NK such evaluations. For our problem, it turns out that such a brute force implementation is not practical on today's PCs, as K needs to be chosen of the order of magnitude of 10^7 for the bound (25) to become sufficiently sharp.

Thus, instead of trying to decide individually whether $C_i \cap Q_j = \emptyset$ for each $Q_j \in \mathcal{Q}$ and $i = 1, \dots, N$, we implement the following divide and conquer algorithm: For each i , initially try to decide for the whole set Θ_0 whether or not $C_i \cap \Theta_0 = \emptyset$. If such a determination can be made, set $J_i = \emptyset$ or $J_i = \{1, 2, \dots, K\}$, respectively. Otherwise, divide Θ_0 into a coarse partition of 4 subsets $\Theta_0 = \{Q_1^c, Q_2^c, Q_3^c, Q_4^c\}$, where each Q_j^c , $j = 1, \dots, 4$ can be covered exactly by a subset of \mathcal{Q} . Try to decide for each Q_j^c whether or not $C_i \cap Q_j^c = \emptyset$. If for any Q_j^c such a determination can be made, include (or not) the appropriate indices in J_i , and do not consider the subspace Q_j^c further. Otherwise, partition Q_j^c again into 4 subsets that can be covered exactly by a subset of \mathcal{Q} , and iterate until either the determination could be made, or the partition is as fine as \mathcal{Q} . The right hand panel of Figure 3 illustrates this approach. The advantage of this algorithm is that it "zooms in" only when necessary. In our implementation, it cuts the number of required

evaluations from 10^7 to around 10^4 to 10^6 , depending on the draw of W_i .

We again relegate the details of the algorithm and the arguments employed for deciding whether or not $C_i \cap Q_j = \emptyset$ to the appendix, where we also discuss the treatment of unbounded subsets $Q_j \in \mathcal{Q}$.

The bound \bar{p} in (25), of course, is still subject to Monte Carlo error, and one might wonder about the effect of the supremum over Θ_0 . Let P^* be the distribution of the N Monte Carlo draws of W_i , $i = 1, \dots, N$.

Lemma 2 For any $x \in \mathbb{R}$, $P^*(\sup_{\theta \in \Theta_0} \hat{p}(\theta) > x) \geq \sup_{\theta \in \Theta_0} P^*(\hat{p}(\theta) > x)$.

Proof. Write E^* for the expectation with respect to P^* . Then, applying Jensen's inequality, $P^*(\sup_{\theta \in \Theta_0} \hat{p}(\theta) > x) = E^*[\mathbf{1}[\sup_{\theta \in \Theta_0} \hat{p}(\theta) > x]] = E^*[\sup_{\theta \in \Theta_0} \mathbf{1}[\hat{p}(\theta) > x]] \geq \sup_{\theta \in \Theta_0} E^*[\mathbf{1}[\hat{p}(\theta) > x]] = \sup_{\theta \in \Theta_0} P^*(\hat{p}(\theta) > x)$. ■

Lemma 2 shows that the P^* distribution of $\sup_{\theta \in \Theta_0} \hat{p}(\theta)$ (weakly) stochastically dominates that of $\hat{p}(\theta)$ for any fixed $\theta \in \Theta_0$. Thus, the P^* probability of observing $\bar{p} \leq 0.05$ is at most as large as the probability of observing $\hat{p}(\theta) \leq 0.05$, even if $\theta \in \Theta_0$ is chosen to maximize actual null rejection probability $E_\theta[\varphi^*(G)]$. Both $\sup_{\theta \in \Theta_0} \hat{p}(\theta)$ and \bar{p} are, in general, upward biased estimates of the size of the test φ^* . A finding of $\bar{p} \leq 0.05$ is thus *stronger* evidence that φ^* is of level 5% compared to any other 5% nominal level test of a single null hypothesis whose critical value was determined by N Monte Carlo draws.

3.4 Power Bound

We now turn to numerical issues that arise in the computation of the power bound via Lemma 1. Without the approximation of integrals by sums as in (21), the Neyman-Pearson test rejects for large values of a statistic of the form

$$\text{LR} = \frac{\int f_\theta dF(\theta)}{\int f_\theta d\Lambda^*(\theta)} = \frac{\int_{0.15}^{0.85} d_0(r) \exp[A_0(r)G(r)^2 + B_0(r)G(r)G(1) + C_0(r)G(1)^2] dr}{\sum_{i=1}^L \int_{a_i}^{b_i} d_i(r) \exp[A_i(r)G(r)^2 + B_i(r)G(r)] dr} \quad (26)$$

where d_i, A_i, B_i and C_i are functions $[0.15, 0.85] \mapsto \mathbb{R}$, and $a_i, b_i \in [0.15, 0.85]$, $i = 1, \dots, L$, and $L, a_i, b_i, d_i, A_i, B_i$, $i = 1, \dots, L$ are numerically determined by the algorithm described in Section 3.1 above. The right hand side in (26) arises from analytically integrating over the conditionally Gaussian distribution of (β, δ) in F of Condition 1 in the numerator, and over a mixture of conditionally Gaussian distributions for δ in the denominator (conditional on ρ). As mentioned above, proper application of Lemma 1 requires the determination of the power of a 5% level test of H_{Λ^*} : The density of G is $\int f_\theta d\Lambda^*(\theta)$ against the alternative H_F : The density of G is $\int f_\theta dF(\theta)$, based on (26).

The usual approach would be to run a Monte Carlo experiment to estimate the critical value and power, using step function approximations to G and Riemann sums instead of integrals in (26). With enough steps (and Monte Carlo draws), this approach delivers any degree of accuracy. However, the test statistic (26) is poorly behaved for this approach: For large breaks (large $|\delta|$), the exponents in (26) take on large values for some r (of the order of magnitude of $\frac{1}{2}\rho\delta^2$), which renders the integrand extremely peaked. Intuitively, for large $|\delta|$, there is ample information about ρ , reflected in a very peaked likelihood. This leads to potentially very poor properties of Riemann sum approximations, casting doubt on the validity of a power bound computed in this way.

For this reason, we develop a lower and upper bound for LR that can be simulated exactly, and use Monte Carlo draws of the lower bound for the computation of the critical value, and Monte Carlo draws from the upper bound for the power computation. This ensures that the resulting upper bound on the power of the test statistic LR is only subject to the usual Monte Carlo error.

The basic element of the bounds is the following inequality: Let $g : [a, b] \mapsto \mathbb{R}$ for some $a > b$ with $\sup_{a \leq x \leq b} |g(x)| < M_g$, and let $\bar{g} = \frac{1}{b-a} \int_a^b g(r) dr$. Then Jensen's inequality and $e^x \leq x \cosh[M_g]$ for all $|x| < M_g$ implies

$$\exp[\bar{g}] \leq \frac{1}{b-a} \int_a^b \exp[g(r)] dr \leq \exp[\bar{g}] \cosh[M_g]. \quad (27)$$

Write any of the integrals in (26) as a sum of integrals over a fine partition of $[0.15; 0.85]$ into J intervals $[c_{j-1}, c_j]$, $j = 1, \dots, J$ with $c_{j-1} < c_j$ for all j and $c_0 = 0.15$ and $c_J = 0.85$. In terms of $W(s) = G(s) - \beta s - \delta \min(\rho, s)$, the representative short integral over $r \in [c_{j-1}, c_j]$ in the denominator has the form

$$\int_{c_{j-1}}^{c_j} d(r) \exp [A(r)W(r)^2 + B(r)W(r)] dr$$

where now $d(r)$, $A(r)$ and $B(r)$ are also functions of $\theta = (\beta, \delta, \rho)$. Using (27), we have

$$\begin{aligned} \underline{d} \exp \left[\underline{A} \overline{W}_j^2 + \underline{B} \bar{W}_j \right] &\leq \frac{1}{c_j - c_{j-1}} \int_{c_{j-1}}^{c_j} d(r) \exp [A(r)W(r)^2 + B(r)W(r)] dr \\ &\leq \bar{d} \exp [\bar{A} \overline{W}_j^2 + \bar{B} \bar{W}_j] \cosh[M_j] \end{aligned} \quad (28)$$

where $\overline{W}_j^2 = \frac{1}{c_j - c_{j-1}} \int_{c_{j-1}}^{c_j} W(r)^2 dr$, $\bar{W}_j = \frac{1}{c_j - c_{j-1}} \int_{c_{j-1}}^{c_j} W(r) dr$, $S_{1,j} = \sup_{c_{j-1} \leq r \leq c_j} |W(r) - \bar{W}_j|$, $S_{2,j} = \sup_{c_{j-1} \leq r \leq c_j} |W(r)^2 - \overline{W}_j^2|$, $M_j = \bar{A} S_{2,j} + \bar{B} S_{1,j}$, and lower and upper bars on d , A and B indicate maximum and minimum values of $d(r)$, $A(r)$ and $B(r)$ on $c_{j-1} \leq r \leq c_j$, respectively, provided $W(r) > 0$ for $c_{j-1} \leq r \leq c_j$. Similar inequalities hold when $W(r) < 0$, when $W(r)$ changes sign on $c_{j-1} \leq r \leq c_j$, and also for the numerator of (26). A lower bound on LR is then

given by replacing integrals in the denominator of (26) with sums over the upper bound in (28) and to replace the integrals in the numerator with sums over corresponding lower bounds, and vice versa for an upper bound.

A Monte Carlo draw from these bounds for LR requires generation of the $4J$ random variables $\{\overline{W}_j^2, \bar{W}_j, S_{1,j}, S_{2,j}\}_{j=1}^J$. Although $\{\bar{W}_j\}_{j=1}^J$ is jointly Gaussian, this is non-trivial, as even the marginal distributions of $\overline{W}_j^2, S_{1,j}$ and $S_{2,j}$ are nonstandard. Fortunately, as detailed in the appendix, it is possible to rely on existing analytical results on the distribution of related statistic of a Wiener process to generate draws $\{\overline{W}_j^{2^l}, \overline{W}_j^{2^u}, \bar{W}_j, S_{1,j}^u, S_{2,j}^u\}_{j=1}^J$ where with very high probability, $\overline{W}_j^{2^l} \leq \overline{W}_j^2 \leq \overline{W}_j^{2^u}$, $S_{1,j} \leq S_{1,j}^u$ and $S_{2,j} \leq S_{2,j}^u$ for all $j = 1, \dots, J$. By amending the above bounds accordingly, this is sufficient for the computation of a Monte Carlo estimate of an upper bound on the power of LR, entirely avoiding any qualifications that arise through the impossibility of generating exact draws of LR under $\int f_\theta d\Lambda^*(\theta)$ and $\int f_\theta dF(\theta)$.

4 Test Statistic and Properties

This section defines the test we suggest for applied work, $\hat{\varphi}^*$, and discusses its properties. In particular, we compare its asymptotic power to the infeasible two-sided t-test with the known break date φ_ρ , and the (approximately) size corrected version of the least-squares break date estimator based test φ_{LS} introduced in Section 2.1.2.

4.1 Definition of Test Statistic

We define the statistic for a general method of moments framework with the $k + 1$ dimensional parameter $\Gamma = (\gamma, \tilde{\gamma})'$, with $\gamma \in \mathbb{R}$ and $\tilde{\gamma} \in \mathbb{R}^k$. The data in a sample of size T is given by $X_T = (x_1, \dots, x_T)$,⁹ and the population moment condition is $E[g(X_t, \Gamma_0)] = 0$ when the true parameter at date t is given by $\Gamma = \Gamma_0$ for some known, \mathbb{R}^m valued function $g(\cdot, \cdot)$. Write $g_t(\Gamma) = g(X_t, \Gamma)$ for notational simplicity. The parameter Γ changes its value from $\Gamma_{pre} = (\gamma_{pre}, \tilde{\gamma}'_{pre})'$ to $\Gamma_{post} = (\gamma_{post}, \tilde{\gamma}'_{post})'$ at some unknown date $\tau \in \mathbb{N}$ in the middle 70% of the sample, $\tau \in [0.15T, 0.85T]$. The hypotheses of interest concern the post break value of the first element γ of Γ ,

$$H_0 : \gamma_{post} = \gamma_{post,0} \quad \text{against} \quad H_1 : \gamma_{post} \neq \gamma_{post,0}. \quad (29)$$

Denote by $\hat{\Gamma}_{pre}(t)$ and $\hat{\Gamma}_{post}(t)$ standard GMM estimators of Γ using data $\{X_s\}_{s=1}^t$ and $\{X_s\}_{s=t+1}^T$, and denote by $\hat{\Omega}_{pre}(t)$ and $\hat{\Omega}_{post}(t)$ the estimators of the covariance matrix of $\hat{\Gamma}_{pre}(t)$ and $\hat{\Gamma}_{post}(t)$,

⁹We allow for double arrays in the data generating process, but omit an explicit dependence on T to ease notation.

respectively. These covariance matrix estimators, as well as the weighting matrix for efficient GMM estimation in the overidentified case, are based on data $\{X_s\}_{s=1}^t$ and $\{X_s\}_{s=t+1}^T$ for the pre and post break estimators, and are consistent for a stable model. When $\{g_t(\Gamma_0)\}$ is serially uncorrelated, this is typically achieved by the usual White form of the covariance matrix estimator, whereas time series correlation in $\{g_t(\Gamma_0)\}$ requires a correction as in Newey and West (1987) or Andrews (1991), for instance.

Our test statistic of (29) only requires evaluation of $\{\hat{\Gamma}_{pre}(t), \hat{\Gamma}_{post}(t), \hat{\Omega}_{pre}(t), \hat{\Omega}_{post}(t)\}$ at the 71 numbers $t = \lfloor lT/100 \rfloor$ for $l = 15, 16, \dots, 85$. Let $\hat{\gamma}_{pre}(l)$ and $\hat{\gamma}_{post}(l)$ be the first element of $\hat{\Gamma}_{pre}(\lfloor lT/100 \rfloor)$ and $\hat{\Gamma}_{post}(\lfloor lT/100 \rfloor)$, and denote by $\hat{\omega}_{pre}^2(l)$ and $\hat{\omega}_{post}^2(l)$ the 1-1 element of $\hat{\Omega}_{pre}(\lfloor lT/100 \rfloor)$ and $\hat{\Omega}_{post}(\lfloor lT/100 \rfloor)$, respectively. Define

$$\begin{aligned}
\widehat{\text{supF}} &= \max_{16 \leq l \leq 85} \frac{(\hat{\gamma}_{post}(l) - \hat{\gamma}_{pre}(l-1))^2}{\hat{\omega}_{post}^2(l) + \hat{\omega}_{pre}^2(l-1)} \\
\hat{\Delta}_{pre}(l) &= l\hat{\gamma}_{pre}(l) - (l-1)\hat{\gamma}_{pre}(l-1), \quad \hat{\Delta}_{post}(l) = (101-l)\hat{\gamma}_{post}(l-1) - (100-l)\hat{\gamma}_{post}(l) \\
\hat{l} &= \arg \min_{16 \leq l \leq 85} \left(\sum_{j=16}^{l-1} \hat{\Delta}_{pre}(j)^2 - (l-1)\hat{\gamma}_{pre}(l-1)^2 + \sum_{j=l+1}^{85} \hat{\Delta}_{post}(j)^2 - (100-l)\hat{\gamma}_{post}(l)^2 \right) \\
\hat{t}_{post} &= \frac{\hat{\gamma}_{post}(\min(\hat{l}+1, 85)) - \gamma_{post,0}}{\sqrt{\hat{\omega}_{post}^2(\min(\hat{l}+1, 85))}} \\
\hat{\omega}^2 &= \frac{(\hat{l}-1)^2}{9900} \hat{\omega}_{pre}^2(\hat{l}-1) + \frac{(100-\hat{l})^2}{9900} \hat{\omega}_{post}^2(\hat{l}) \\
\widehat{\text{LR}} &= \frac{\sum_{l=15}^{85} \frac{\exp[\frac{1}{2} \frac{\sigma_{pre}^2(\hat{\gamma}_{pre}(l) - \gamma_{post,0})^2 l^2}{100^2 \hat{\omega}^2 v(l, \sigma_{pre}^2)} + \frac{1}{2} \frac{\sigma_{\beta}^2(\hat{\gamma}_{post}(l) - \gamma_{post,0})^2 (100-l)^2}{100^2 \hat{\omega}^2 v(100-l, \sigma_{\beta}^2)}]}{71 \sqrt{v(l, \sigma_{pre}^2) v(100-l, \sigma_{\beta}^2)}}}{\sum_{j=1}^{18} \sum_{l=a_j}^{b_j} \frac{p_j \exp[-\frac{1}{2} \frac{\mu_{\delta,j}^2 l}{100 v(l, \sigma_{\delta,j}^2)} + \frac{1}{2} \frac{\sigma_{\delta,j}^2(\hat{\gamma}_{pre}(l) - \gamma_{post,0})^2 l^2}{100^2 \hat{\omega}^2 v(l, \sigma_{\delta,j}^2)}] \cosh[\frac{(\hat{\gamma}_{pre}(l) - \gamma_{post,0}) \mu_{j,l}}{100 v(l, \sigma_{\delta,j}^2) \hat{\omega}}]}{(b_j - a_j + 1) \sqrt{v(l, \sigma_{\delta,j}^2)}}}
\end{aligned} \tag{30}$$

where $v(l, \sigma^2) = 1 + \sigma^2 l/100$, $\sigma_{pre}^2 = 378$, $\sigma_{\beta}^2 = 22$, and a_j, b_j, p_j, μ_j and $\sigma_{\delta,j}^2$ are defined in Table 2. The test $\hat{\varphi}^*$ of (29) rejects H_0 if and only if either (i) $\widehat{\text{supF}} > 90$ and $|\hat{t}_{post}| > 2.01$, or (ii) $\widehat{\text{supF}} \leq 90$ and $\widehat{\text{LR}} > 2.41$.¹⁰

The intuition for these computations is roughly as follows. The statistic $\widehat{\text{supF}}$ is the largest F-statistic of the null hypothesis that the value of γ in the first $(l-1)\%$ is equal to the value of γ in the last $(100-l)\%$, maximized over $16 \leq l \leq 85$. By leaving out the middle 1% of observations, it is ensured that for any true break fraction within the middle 70% of the sample, one of these F-statistics (often, the largest) only involves estimators $(\hat{\gamma}_{pre}(l), \hat{\omega}_{pre}^2(l))$ and $(\hat{\gamma}_{post}(l), \hat{\omega}_{post}^2(l))$ from stable periods. When $\widehat{\text{supF}} > 90$, that is, when there is strong

¹⁰One obtains the test φ^* of Proposition 1 and defined in the appendix by setting $\hat{\gamma}_{pre}(l) = 100G(l/100)/l$, $\hat{\gamma}_{post}(l) = (G(1) - G(l/100))/(1 - l/100)$, $\hat{\omega}_{pre}^2(l) = 100/l$ and $\hat{\omega}_{post}^2(l) = 100/(100-l)$.

Table 2: Weights, Intervals, Means and Variances of the Approximate Least Favorable Distribution

| | | | | | | | | | |
|--------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| j | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| p_j | 0.588 | 0.123 | 0.067 | 0.057 | 0.038 | 0.032 | 0.026 | 0.02 | 0.009 |
| a_j | 0.15 | 0.85 | 0.85 | 0.2 | 0.75 | 0.2 | 0.2 | 0.75 | 0.45 |
| b_j | 0.85 | 0.85 | 0.85 | 0.74 | 0.85 | 0.74 | 0.74 | 0.82 | 0.59 |
| σ_j^2 | 100 | 10 | 4 | 300 | 200 | 10 | 3 | 10 | 10 |
| μ_j | 20 | 5 | 3 | 16 | 28 | 9 | 6 | 7 | 11 |
| j | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 |
| p_j | 0.009 | 0.008 | 0.006 | 0.005 | 0.004 | 0.004 | 0.002 | 0.001 | 0.001 |
| a_j | 0.7 | 0.15 | 0.15 | 0.6 | 0.8 | 0.6 | 0.83 | 0.85 | 0.75 |
| b_j | 0.74 | 0.19 | 0.24 | 0.69 | 0.82 | 0.69 | 0.84 | 0.85 | 0.82 |
| σ_j^2 | 10 | 10 | 200 | 10 | 10 | 3 | 10 | 3 | 3 |
| μ_j | 9 | 5 | 28 | 12 | 11 | 8 | 13 | 15.5 | 13 |

evidence for the occurrence of a break, the test $\hat{\varphi}^*$ rejects if the usual t-statistic \hat{t}_{post} is larger than 2.01 (rather than the usual 1.96) in absolute value, where \hat{t}_{post} uses \hat{l} to determine the appropriate post-break data.

In (approximately) linear and stationary models, $\hat{\gamma}_{pre}(l)$ is centred around the average parameter value of the first $l\%$ of the sample, $[lT/100]^{-1} \sum_{t=1}^{[lT/100]} (\gamma_{pre} + (\gamma_{post} - \gamma_{pre})\mathbf{1}[t > \tau])$, and similarly, $\hat{\gamma}_{post}(l)$ is centred about the average parameter value of the last $(100 - l)\%$ of the sample. The statistics $\hat{\Delta}_{pre}(l)$ and $\hat{\Delta}_{post}(l)$ thus approximately estimate the value of the parameter in the l th percent of the sample. If the true break τ is in the l_0 th percent of the sample, then $\hat{\Delta}_{pre}(l)$ should be approximately equal to $\hat{\gamma}_{pre}(l_0)$ for all $l < l_0$, and similarly for $\hat{\Delta}_{post}(l)$. The break fraction estimator \hat{l} is determined by minimizing a corresponding least squares criterion, again dropping the middle one percent. By adding one to \hat{l} , it is ensured that with high probability, $\hat{l} + 1$ is at least as large as the true break fraction in percentage points, so that \hat{t}_{post} is based on estimates from a stable post-break period. The advantage of the least squares criterion (30) based on $(\hat{\Delta}_{pre}(l), \hat{\Delta}_{post}(l))$ over, say, an analogue to the F-statistics that underlie $\widehat{\text{supF}}$, is that \hat{l} in (30) has appealing properties under relatively weak conditions also when the break magnitude is not small—cf. Proposition 4 below.

In absence of strong evidence for a break, that is $\widehat{\text{supF}} \leq 90$, the test $\hat{\varphi}^*$ switches to deciding the null hypothesis based on a likelihood ratio statistic. The numerator of this statistic is the result of the weighting over alternative values for γ_{post} , break dates and break magnitudes

of Condition 1. The denominator is the result of a weighting over break dates and break magnitudes that make detection of this alternative as difficult as possible, the numerically determined approximate least favorable distribution discussed in Section 3, ensuring size control of $\hat{\varphi}^*$ even when information about the true break date is scarce. In this likelihood ratio statistic, the scale of $\hat{\gamma}_{pre}(l)$ and $\hat{\gamma}_{post}(l)$ is normalized by $\hat{\omega}$. If the break data τ satisfies $(\hat{l} - 1)/100 \leq \tau/T \leq \hat{l}/100$, then the estimator $\hat{\omega}^2$ is based on variance estimators of stable models, which improves the small sample properties of $\widehat{\text{LR}}$.

4.2 Asymptotic Properties

Proposition 4 (i) Suppose $T^{1/2}(\gamma_{pre} - \gamma_{post}) = \omega\delta \in \mathbb{R}$, $\tau = \lfloor \rho T \rfloor$ with $\rho \in [0.15, 0.85]$ and

$$\begin{aligned} & \{T^{1/2}(\hat{\gamma}_{pre}(l) - \gamma_{post}), T^{1/2}(\hat{\gamma}_{post}(l) - \gamma_{post}), T\hat{\omega}_{pre}^2(l), T\hat{\omega}_{post}^2(l)\}_{l=15}^{85} \\ & \Rightarrow \left\{ \omega \frac{G_0(l/100)}{l/100}, \omega \frac{G_0(1) - G_0(l/100)}{1 - l/100}, \frac{\omega^2}{l/100}, \frac{\omega^2}{1 - l/100} \right\}_{l=15}^{85} \end{aligned} \quad (31)$$

where $G_0(s) = W(s) + \delta \min(\rho, s)$, and $\omega > 0$. (i.a) If $T^{1/2}(\gamma_{post} - \gamma_{post,0}) = \omega\beta \in \mathbb{R}$, then $\hat{\varphi}^*(X_T) \Rightarrow \varphi^*(G)$. (i.b) If $T^{1/2}(\gamma_{post} - \gamma_{post,0}) \rightarrow \pm\infty$, then $\hat{\varphi}^*(Y_T) \xrightarrow{p} 1$.

(ii) Suppose $T^{1/2}(\gamma_{pre} - \gamma_{post}) \rightarrow \pm\infty$, $\tau = \lfloor \rho T \rfloor$ with $\rho \in [0.15, 0.85]$, and, for some $\omega_{pre}, \omega_{post} > 0$,

$$\begin{aligned} & \{T^{1/2}(\hat{\gamma}_{pre}(l) - \gamma_{pre}), T\hat{\omega}_{pre}^2(l)\}_{l=15}^{\lfloor 100\rho \rfloor} \Rightarrow \left\{ \omega_{pre} \frac{G_0(l/100)}{l/100}, \frac{\omega_{pre}^2}{l/100} \right\}_{l=15}^{\lfloor 100\rho \rfloor} \\ & \{T^{1/2}(\hat{\gamma}_{post}(l) - \gamma_{post}), T\hat{\omega}_{post}^2(l)\}_{l=\lceil 100\rho \rceil}^{85} \Rightarrow \left\{ \omega_{post} \frac{G_0(1) - G_0(l/100)}{1 - l/100}, \frac{\omega_{post}^2}{1 - l/100} \right\}_{l=\lceil 100\rho \rceil}^{85} \\ & P(T^{1/2}(\hat{\gamma}_{pre}(\lfloor 100\rho \rfloor + 1) - \gamma_{pre}) > M) \rightarrow 1, P(T^{1/2}(\hat{\gamma}_{post}(\lceil 100\rho \rceil - 1) - \gamma_{post}) > M) \rightarrow 1 \end{aligned} \quad (32)$$

for any $M \in \mathbb{R}$, where $\lceil x \rceil = -\lfloor -x \rfloor$. (ii.a) If $\gamma_{post} = \gamma_{post,0}$, then $\limsup_{T \rightarrow \infty} E[\hat{\varphi}^*(X_T)] \leq 5\%$. (ii.b) If $T^{1/2}(\gamma_{post} - \gamma_{post,0}) = \omega_{post}\beta \in \mathbb{R}$ and $\rho \notin R_{grid} = \{0.15, 0.16, \dots, 0.84\}$, then $\lim_{T \rightarrow \infty} \hat{\varphi}^*(X_T) = \mathbf{1}[|Z + \beta\sqrt{0.98 - \lfloor 100\rho \rfloor / 100}| > 2.01]$. (iii.c) If $T^{1/2}(\gamma_{post} - \gamma_{post,0}) \rightarrow \pm\infty$, then $\hat{\varphi}^*(X_T) \xrightarrow{p} 1$.

(iii) Suppose $\gamma_{post} = \gamma_{post,0}$; (31) holds for all sequences $(\rho_T, \delta_T) = (\tau/T, T^{1/2}(\gamma_{pre} - \gamma_{post})) \rightarrow (\rho, \delta) \in [0.15, 0.85] \times \mathbb{R}$; (32) holds for all sequences $(\rho_T, \delta_T) \rightarrow (\rho, \delta) \in ([0.15, 0.85] \setminus R_{grid}) \times \{-\infty, +\infty\}$; and for all for all sequences $(\rho_T, \delta_T) \rightarrow (\rho, \delta) \in R_{grid} \times \{-\infty, +\infty\}$, there exists a subsequence $(\rho_{T'}, \delta_{T'}) \rightarrow (\rho, \delta)$ such that along this subsequence, (32) holds except for

$$\begin{aligned} & (T^{1/2}(\hat{\gamma}_{pre}(l_0) - \gamma_{pre}), T\hat{\omega}_{pre}^2(l_0)) \Rightarrow (\omega_{pre} \frac{G_0(l_0/100)}{l_0/100} + \nu_{pre}(l_0), \frac{\omega_{pre}^2}{l/100}) \\ & (T^{1/2}(\hat{\gamma}_{post}(l_0) - \gamma_{pre}), T\hat{\omega}_{pre}^2(l_0)) \Rightarrow (\omega_{post} \frac{G_0(1) - G_0(l_0/100)}{1 - l_0/100} + \nu_{post}(l_0), \frac{\omega_{post}^2}{1 - l_0/100}) \end{aligned} \quad (33)$$

where $l_0 = 100\rho$ and $\nu_{pre}(l_0), \nu_{post}(l_0) \in \mathbb{R} \cup \{-\infty, +\infty\}$ are nonrandom constants, of which at most one is nonzero. Then $\limsup_{T \rightarrow \infty} \sup_{\theta \in \Theta_0} E_\theta[\hat{\varphi}^*(X_T)] \leq \alpha^* \simeq 5\%$.

Part (i) of Proposition 4 considers the case where the pre and post break value of γ only differ of the order $T^{-1/2}$. In that neighborhood, the GMM estimators do not contain enough information to pin down the true break fraction exactly, even asymptotically. This asymptotic embedding represents the small sample problem with substantial uncertainty about the true break date. The proposition establishes that for local alternatives, where correspondingly, the true value γ_{post} differs by the order $T^{-1/2}$ from the hypothesized value $\gamma_{post,0}$, the asymptotic properties of $\hat{\varphi}^*$ are just like those of φ^* in the limiting problem discussed in Section 2.1. So in particular, by Proposition 1, the test $\hat{\varphi}^*$ has asymptotic level $\alpha^* \simeq 5\%$, and it has asymptotic weighted average power under Condition 1 of $\pi^* \simeq 49\%$. Furthermore, $\hat{\varphi}^*$ is consistent in this scenario against any alternative where γ_{post} is outside the $T^{-1/2}$ neighborhood of the hypothesized value.

These results rely on the high level condition (31). The sequence of statistics $\{\hat{\gamma}_{pre}(l), \hat{\gamma}_{post}(l), \hat{\omega}_{pre}(l), \hat{\omega}_{post}(l)\}$ are a special case of partial sample GMM estimators analyzed by Andrews (1993). In particular, his primitive Assumption 1 with an appropriate modification to account for the local break in the parameter value imply (31)—see the appendix for details. Alternatively, the approach of Li and Müller (2007) could be amended to yield (31) under a different set of assumptions. Also, for the special case of maximum likelihood estimation, Condition 2 in the appendix can be shown to imply (31) for both the average Hessian and outer product of scores covariance estimators $\hat{\Omega}_{pre}$ and $\hat{\Omega}_{post}$. The conclusions of part (i) thus hold for a wide range of models.¹¹

Part (ii) of Proposition 4 establishes the asymptotic properties of $\hat{\varphi}^*$ when the break is (very) large relative to the sample information about γ . The test again controls size and is consistent against any non-local alternative. Under local alternatives $T^{1/2}(\gamma_{post} - \gamma_{post,0}) = \beta \neq 0$, it asymptotically corresponds to a two-sided t-test about unit variance normal with non-centrality parameter of $\sqrt{0.98 - [100\rho]/100}$ and critical value 2.01, at least as long $\rho \neq R_{grid}$.¹² In comparison, the two-sided 5% level t-test based on post break data (assuming the break date was known) converges to $\varphi_\rho(G) = \mathbf{1}[|\frac{G(1)-G(\rho)}{\sqrt{1-\rho}}| > 1.96]$, and thus has a non-centrality parameter of $\beta\sqrt{1-\rho}$. With $\beta \sim \mathcal{N}(0, 22)$ as in Condition 1, the average asymptotic power loss of $\hat{\varphi}^*$ relative

¹¹As pointed out by Hansen (2000), most tests for parameter stability do not controls asymptotic size under 'global' heterogeneity of average variance across different parts of the sample. This also applies to $\hat{\varphi}^*$, at least for local breaks. However, by picking the 71 grid points in the definition of $\hat{\varphi}^*$ not relative to the sample proportion, but relative to a suitably chosen cumulative information (that typically would be need to be estimated with sufficient precision), one could induce the convergence (31) even for models with 'global' variance heterogeneity.

¹²This qualification is necessary because if $\rho \in R_{grid}$, \hat{l} potentially takes on the two values $[100\rho] + 1$ and $[100\rho] + 2$ with positive probability, even asymptotically, despite the diverging break magnitude $T^{1/2}(\gamma_{pre} - \gamma_{post})$.

to φ_ρ over $\rho \in [0.15, 0.85]$ is approximately 1.7 percentage points, with a largest difference of 3.6 percentage points occurring at $\rho \rightarrow 0.83$.

The results in part (ii) of Proposition 4 require two sets of assumptions. On the one hand, the statistics $\{\hat{\gamma}_{pre}(l), \hat{\omega}_{pre}^2(l)\}$ and $\{\hat{\gamma}_{post}(l), \hat{\omega}_{post}^2(l)\}$ have to behave in the usual way *over the stable pre and post break periods*, respectively, and be asymptotically independent. Note that the limiting variance may change at the parameter break date; this accommodates, say, changes in the variance of the AR(1) coefficient estimator that are induced by a break in the AR(1) coefficient. One can again invoke the primitive conditions of Andrews (1993) to justify these convergences. On the other hand, the estimators $\hat{\gamma}_{pre}(l)$ and $\hat{\gamma}_{post}(l)$ must diverge from the $T^{-1/2}$ neighborhood of the pre and post parameter values γ_{pre} and γ_{post} for two values of l that involve a positive fraction of post and pre break data, respectively.¹³ For a non-local but shrinking break magnitude $|\gamma_{pre} - \gamma_{post}| \rightarrow 0$ and $T^{1/2}(\gamma_{pre} - \gamma_{post}) \rightarrow \pm\infty$, $\hat{\gamma}_{pre}(l)$ typically estimates the average parameter value in the first $l\%$ of the sample, that is

$$\frac{\hat{\gamma}_{pre}(l) - \gamma_{pre}}{\gamma_{post} - \gamma_{pre}} \xrightarrow{p} \mathbf{1}[l > \lfloor 100\rho \rfloor] \frac{l - \lfloor 100\rho \rfloor}{l} \neq 0, \quad (34)$$

and similarly for $\hat{\gamma}_{post}(l)$, which is clearly sufficient for the purposes of Proposition 4 (ii). The convergence (34) can be shown to hold, for instance, under the high-level Condition 1 of Li and Müller (2007) by proceeding as in their Theorem 1. For a fixed break magnitude $\gamma_{post} - \gamma_{pre}$, the distortionary effects of the break becomes even stronger, of course, and barring pathological cancellations, one would expect the condition in Proposition (ii) to be satisfied. We refrain from a detailed discussion of sufficient primitive conditions for the sake of brevity.

Finally, part (iii) of Proposition 4 shows that the test $\hat{\varphi}^*$ controls size uniformly in large samples—for T large enough, no choice of $\theta \in \Theta_0$ leads to a rejection probability substantially above 5%. For local breaks $\gamma_{pre} - \gamma_{post} = O(T^{-1/2})$, the assumption is only that the convergences in part (i) hold for all converging sequences $(\rho_T, \delta_T) \rightarrow (\rho, \delta)$. We show in the appendix that the primitive assumptions of Andrews (1993) are again sufficient. For non-local breaks, care must be taken for the case $\rho_T \rightarrow \rho \in R_{grid}$. If $\rho_T \rightarrow \rho \in R_{grid}$ with $\rho_T < \rho$, then $\hat{\gamma}_{pre}(100\rho)$ is an estimator from an unstable model, with a shrinking fraction of the data stemming from the post break model. Depending on the rate at which $\delta_T \rightarrow \infty$, this contamination shifts the center of the asymptotic distribution of $\hat{\gamma}_{pre}(100\rho)$ by $\nu_{pre}(100\rho)$ —typically, given the arguments above (34), one would expect $\nu_{pre}(100\rho) \in \mathbb{R}$ if $\delta_T(\rho_T - \rho) \rightarrow \xi \in \mathbb{R}$, and it is sufficient for the last assumption in Proposition 4 (iii) to assume that when $(\rho_T, \delta_T) \rightarrow (\rho, \delta) \in R_{grid} \times \{-\infty + \infty\}$,

¹³Note, however, that nothing is assumed about the behavior of the variance estimators $\{\hat{\omega}_{pre}(l), \hat{\omega}_{post}(l)\}$ that involve subsets of unstable periods.

(32) holds except for (33) for all sequences $\delta_T(\rho_T - \rho) \rightarrow \xi \in \mathbb{R} \cup \{-\infty, +\infty\}$.¹⁴

By Proposition 4 part (i), the asymptotic local power of $\hat{\varphi}^*$ equals the power of φ^* . As discussed in Section 2.2, at least in linear regressions with a single coefficient, size corrected 5% level inference based on the least squares break date estimator (where the critical value of the t-statistic is 4.9 instead of 1.96) has asymptotic power equal to the power of $\varphi_{\text{LS}}(G) = \mathbf{1}[|G(1) - G(\hat{\rho})|/\sqrt{1 - \hat{\rho}} > 4.9]$, where $\hat{\rho} = \operatorname{argmax}_{r \in [0.15; 0.85]} (G(r) - rG(1))^2 / (r(1 - r))$, under suitable conditions. Finally, under the conditions of Proposition 4 part (i), the infeasible test using actual post break data only yields asymptotic local power equal to the power of $\varphi_\rho(G) = \mathbf{1}[|G(1) - G(\rho)|/\sqrt{1 - \rho} > 1.96]$, and the feasible test that uses only the last 15% of the data has asymptotic local power equal to $\varphi_{0.85}(G) = \mathbf{1}[|G(1) - G(0.85)|/\sqrt{0.15} > 1.96]$. Figure 4 compares the power of these tests for selected values of ρ and δ . All tests are invariant under the transformation $G \rightarrow -G$, so there is not need to plot the the power for negative δ . As one might expect given the large critical value of 4.9, φ_{LS} has very low power. In contrast, φ^* is almost always more powerful than $\varphi_{0.85}$, often substantially so, and comes close to the power of the infeasible benchmark φ_ρ for large $|\delta|$.

We also investigated asymptotic size control of $\hat{\varphi}^*$ in models with a local break where the transition to the post break parameter value is smooth rather than a sudden shift at ρ . In particular, we considered parameter paths of the form $T^{1/2}(\gamma_t - \gamma_{post,0}) = \delta(\Phi_{0.05}(t/T - \rho) - 1)$, where $\Phi_{0.05}$ is the cdf of a mean zero normal with standard deviation 0.05. For large $|\delta|$, the test is very substantially oversized, but for $|\delta| < 15$, the asymptotic rejection probability of $\hat{\varphi}^*$ is always below 7%, at least for $\rho \in \{0.25, 0.5, 0.75\}$. For most moderately sized breaks encountered in practice, these results suggest that $\hat{\varphi}^*$ continuous to yield approximately valid inference about the post break parameter value even if the transition to the post break value takes place over a nonnegligible fraction of the sample.

4.3 Small Sample Properties

We now turn to the small sample properties of the test $\hat{\varphi}^*$ suggested here, and compare it to the infeasible test $\hat{\varphi}_\rho$ that corresponds to standard inference using actual post break data only. We consider two Monte Carlo designs: Inference about the post break mean of an otherwise

¹⁴Inspection of the proof of Proposition 4 shows that these assumptions are only needed to ensure that $(\{\hat{\gamma}_{pre}(l)\}_{l=1}^{l_0}, \hat{\Delta}_{post}(l_0 + 1))$ are asymptotically independent of $(\{\hat{\gamma}_{post}(l), \hat{\omega}_{post}^2(l)\}_{l=l_0+1}^{l_0+2})$, as this independence is enough to establish that the mixture of the two \hat{t}_{post} statistics with $\hat{l} = l_0 + 1$ and $\hat{l} = l_0 + 2$ controls size. The condition could dispensed with entirely by increasing the critical value of \hat{t}_{post} to 2.07, since $P(|Z_1| < 2.07 \text{ and } |Z_2| < 2.07) > 0.95$ for all bivariate normal Z_1, Z_2 with $Z_1 \sim Z_2 \sim \mathcal{N}(0, 1)$ and correlation of at least $\sqrt{15/16}$, so that no mixture of the two \hat{t}_{post} -statistics can induce overrejections.




Figure 4: Asymptotic power of approximately optimal test φ^* and of the least-squared date estimator based test φ_{LS} compared to the infeasible test φ_ρ with break date known, and the test using only the last 15% of the observations $\varphi_{0.85}$

stationary scalar time series, and inference about the post break value of the moving average parameter of a scalar time series.

For inference about the mean, we consider a series of $T = 180$ observations (think of 45 years of quarterly data) $\{y_t\}_{t=1}^T$, where

$$y_t = \beta + \delta T^{-1/2} \mathbf{1}[t \leq \rho T] + u_t$$

and the zero mean disturbance u_t is either i.i.d. standard normal, or follows a stationary Gaussian AR(1) with coefficient 0.4, or follows a (positively autocorrelated) stationary MA(1) with coefficient 0.4. In all cases, we estimate the variance of the sample means by the quadratic spectral long run variance estimator with automatic bandwidth selection based on an AR(1) model, as suggested in Andrews (1991). Table 3 shows size and power of $\hat{\varphi}^*$ in this scenario, along with the properties of $\hat{\varphi}_\rho$ (the usual t-test using only actual post break data), and of $\hat{\varphi}_{0.85}$ (the usual t-test using only the last 15% of the data). With autocorrelated disturbances u_t , there are some size distortions for $\hat{\varphi}^*$, although they are smaller than those of the infeasible benchmark statistic $\hat{\varphi}_\rho$. In comparison, unreported simulations show that (non-size corrected) inference based on the least squares break date estimator is subject to large size distortions; in the MA(1) case, for instance, the null rejection probability ranges from 8.1% to 22.5% for the values of ρ and δ considered in Table 3. The (not size corrected) power results of Table 3 at least qualitatively correspond closely to the asymptotic results in Figure 4.

In the MA(1) design, we generate data from the model

$$y_t = \mu + \varepsilon_t + \mathbf{1}[t \leq \rho T] \psi_{pre} \varepsilon_{t-1} + \mathbf{1}[t > \rho T] \psi_{post} \varepsilon_{t-1}$$

with $\varepsilon_t \sim \text{i.i.d. } \mathcal{N}(0, \sigma^2)$ and $T = 480$ (think of 40 years of monthly data). We test the hypothesis $H_0 : \psi_{post} = 0$, which amounts to testing whether y_t is currently forecastable. Table 4 shows the null and alternative rejection probabilities for this model, where the parameters are estimated by (unconditional) maximum likelihood. Size distortions are mostly larger for $\hat{\varphi}^*$ than for $\hat{\varphi}_\rho$, but remain modest for most considered parameter values. We experimented with smaller sample sizes and found worse size control, sometimes substantially so. Good properties of $\hat{\varphi}^*$ rely on reasonable accuracy of the usual large sample approximations over all partial sample estimators. But with small sample sizes T , the estimators over the first and last 15% are based on very few observations, which leads to well documented problems with the MA(1) maximum likelihood estimator.

Table 3: Small Sample Size and Power of Suggested Test in Location Model

| ρ | Size ($\beta = 0$) | | | | | Power ($\beta = 4$) | | | | |
|---------------------|-----------------------------|-----|-----|------|----------------------|-----------------------------|------|------|------|----------------------|
| | $\hat{\varphi}^*, \delta =$ | | | | $\hat{\varphi}_\rho$ | $\hat{\varphi}^*, \delta =$ | | | | $\hat{\varphi}_\rho$ |
| | 1 | 4 | 8 | 16 | | 1 | 4 | 8 | 16 | |
| i.i.d. disturbances | | | | | | | | | | |
| 0.25 | 4.9 | 5.1 | 4.8 | 4.2 | 5.3 | 41.6 | 53.2 | 78.2 | 89.8 | 93.2 |
| 0.50 | 4.8 | 5.1 | 5.2 | 4.8 | 5.7 | 41.3 | 50.7 | 68.0 | 74.6 | 80.7 |
| 0.75 | 4.7 | 4.5 | 5.6 | 5.9 | 6.0 | 41.6 | 42.0 | 44.6 | 44.2 | 52.7 |
| AR(1) disturbances | | | | | | | | | | |
| 0.25 | 8.9 | 9.0 | 8.1 | 7.4 | 8.2 | 52.2 | 63.8 | 83.2 | 92.6 | 94.6 |
| 0.50 | 9.1 | 8.7 | 8.4 | 7.9 | 9.0 | 51.5 | 60.5 | 74.4 | 80.1 | 85.0 |
| 0.75 | 9.0 | 7.9 | 8.6 | 10.3 | 11.6 | 51.1 | 49.6 | 52.5 | 55.3 | 63.1 |
| MA(1) disturbances | | | | | | | | | | |
| 0.25 | 7.0 | 7.3 | 6.4 | 5.8 | 6.9 | 48.0 | 59.5 | 80.9 | 91.2 | 93.6 |
| 0.50 | 7.1 | 7.2 | 7.0 | 6.3 | 7.3 | 47.5 | 56.3 | 71.4 | 77.8 | 82.5 |
| 0.75 | 6.8 | 7.0 | 7.7 | 8.3 | 8.9 | 47.5 | 46.6 | 48.7 | 50.0 | 58.1 |

Table 4: Small Sample Size and Power of Suggested Test in MA(1) Model

| ρ | Size ($\psi_{post} = 0$) | | | | | Power ($\psi_{post} = 0.2$) | | | | |
|--------|---------------------------------|-----|------|------|----------------------|---------------------------------|------|------|------|----------------------|
| | $\hat{\varphi}^*, \psi_{pre} =$ | | | | $\hat{\varphi}_\rho$ | $\hat{\varphi}^*, \psi_{pre} =$ | | | | $\hat{\varphi}_\rho$ |
| | 0.1 | 0.3 | 0.5 | 0.7 | | -0.1 | 0.1 | 0.3 | 0.5 | |
| 0.25 | 5.7 | 5.6 | 5.0 | 5.2 | 5.2 | 60.3 | 85.5 | 96.9 | 96.9 | 97.5 |
| 0.50 | 6.0 | 7.1 | 8.4 | 8.3 | 5.9 | 61.3 | 85.4 | 88.6 | 89.1 | 89.7 |
| 0.75 | 5.7 | 8.9 | 12.0 | 13.3 | 7.0 | 53.6 | 51.0 | 56.7 | 66.6 | 64.4 |

5 Conclusion

Models with discrete breaks in the parameters at an unknown or uncertain date have become popular in empirical work in recent years. This paper shows that inference about pre and post break parameters using the estimated break date as the true break date leads to substantially oversized tests and confidence intervals as long as the break magnitude is not very large relative to the sampling uncertainty about the parameters. For the important special case of a single break at an unknown date and a single parameter of interest, we derive an alternative test with uniform asymptotic size control that demonstrably comes close to maximizing a weighted average power criterion.

While the test is entirely straightforward to apply and not particularly computationally burdensome, the test statistic is certainly not particularly elegant. The reason is that our derivation of the approximately power maximizing test does not rely on a particular 'trick' to deal with the nuisance parameter under the null hypothesis, unlike, say, Jansson and Moreira (2006) or Andrews, Moreira, and Stock (2006), for the simple reason that no such trick seems to be applicable. The strength of our approach to identifying a suitable test is precisely its generic nature. The suggested algorithm is computationally intensive and heavily involves numerical approximations, but given our algorithm to check size control, its output is as valid a test as any whose critical value is determined by Monte Carlo simulation.

In many ways, the problem of pre and post break parameter inference has a number of features—such as the very peaked likelihood for large breaks, the two dimensions in the nuisance parameter space and the dependence on an infinite dimensional disturbance—that arguably make the engineering challenges particularly severe. We would therefore expect that many aspects of the approach here could be successfully applied in other testing problems with nuisance parameters under the null hypothesis that cannot be estimated consistently. For instance, in the context of inference in models with parameters undergoing a single break, one might consider inference about the magnitude of the change (leading to the break date as the single nuisance parameter after invoking invariance to eliminate the average value of the parameter), or optimal inference about the break date (leaving the break magnitude as the only relevant nuisance parameter after invoking the same invariance), or the same question considered in the present paper, but in the low frequency framework recently advanced by Müller and Watson (2008b). The one clearly binding constraint in the applicability of the approach is the dimension of the nuisance parameter space, as it is unclear how one could numerically check whether a given test controls size for all values of a high dimensional nuisance parameter.

6 Appendix

Definition of φ^* :

$\varphi^*(G) = \mathbf{1}[\text{supF}_0 > 90]\mathbf{1}[\hat{t}_0 > 2.00] + \mathbf{1}[\text{supF}_0 \leq 90]\mathbf{1}[\text{LR}_0 > 2.56]$, where

$$\begin{aligned} \text{supF}_0 &= \max_{16 \leq l \leq 85} \frac{100 \left((100-l)G\left(\frac{l-1}{100}\right) - (l-1)(G(1) - G\left(\frac{l}{100}\right)) \right)^2}{99(100-l)(l-1)} \\ S_0(l) &= \frac{100(100G\left(\frac{l}{100}\right) - lG(1))^2}{l(100-l)} + 100^2 \frac{(lG\left(\frac{l-1}{100}\right) - (l-1)G\left(\frac{l}{100}\right))^2}{l(l-1)} \\ \hat{l}_0 &= \arg \max_{16 \leq l \leq 85} S_0(l) \\ \hat{t}_0 &= \frac{G(1) - G(\min(\hat{l}_0 + 1, 85)/100)}{\sqrt{1 - \min(\hat{l}_0 + 1, 85)/100}} \\ \text{LR}_0 &= \frac{\sum_{l=15}^{85} v(l, \sigma_{pre}^2)^{-1/2} v(100-l, \sigma_\beta^2)^{-1/2} \exp\left[\frac{1}{2} \frac{\sigma_{pre}^2 G(l/100)^2}{v(l, \sigma_{pre}^2)} + \frac{1}{2} \frac{\sigma_\beta^2 (G(1) - G(l/100))^2}{v(100-l, \sigma_\beta^2)}\right]}{\sum_{j=1}^{18} \sum_{l=a_j}^{b_j} \frac{p_j}{b_j - a_j + 1} v(l, \sigma_{\delta,j}^2)^{-1/2} \exp\left[-\frac{1}{2} \frac{\mu_{\delta,j}^2 l}{100v(l, \sigma_{\delta,j}^2)} + \frac{1}{2} \frac{\sigma_{\delta,j}^2 G(l/100)^2}{v(l, \sigma_{\delta,j}^2)}\right] \cosh\left[\frac{\mu_j G(l/100)}{v(l, \sigma_{\delta,j}^2)}\right]} \end{aligned}$$

and $v(l, \sigma^2) = 1 + \sigma^2 l/100$, $\sigma_{pre}^2 = 378$, $\sigma_\beta^2 = 22$, and $p_j, \sigma_{\delta,j}^2, \mu_j$ are defined in Table 2.

Proof of Proposition 2:

By Girsanov's Theorem, the log of the Radon-Nikodym (RN) derivative of the distribution of $(G, \tilde{G})'$ with respect to the distribution of $(W, (A_{21}W + A_{22}\tilde{W}_k)')$ is the r.h.s. of (18).

(i) Fix arbitrary values $\rho_0 \in [0.15, 0.85]$, $\delta_0 \in \mathbb{R}$, $\tilde{\beta}_0, \tilde{\delta}_0 \in \mathbb{R}^k$, and consider the construction of the most powerful unbiased test against the specific alternative $H_1^s(b) : \beta = b, \delta = \delta_0 - b, \tilde{\beta} = \tilde{\beta}_0 + bA_{21}, \tilde{\delta} = \tilde{\delta}_0 - bA_{21}$ $\rho = \rho_0$, which is indexed by $b \in \mathbb{R}$. Any 5% level test for the unrestricted null hypothesis is also of level 5% under the specific null hypothesis $H_0 : \beta = 0, \delta = \delta_0, \tilde{\beta} = \tilde{\beta}_0, \tilde{\delta} = \tilde{\delta}_0, \rho = \rho_0$. The log of the RN derivative of the distribution of $(G, \tilde{G})'$ under $H_1^s(b)$ with respect to the distribution under H_0^s then becomes $b(G(1) - G(\rho_0)) - \frac{1}{2}(1 - \rho_0)b^2$. The experiment of testing H_0^s against H_1^s based on the observation $(G, \tilde{G})'$ is therefore *equivalent* (in the sense of equivalence of statistical experiments) to inference about b in the Gaussian shift experiment of observing $X \sim \mathcal{N}(b, 1 - \rho_0)$. In particular, for any test of H_0^s against $H_1^s(b)$, there exists a test $\varphi_X : \mathbb{R} \mapsto [0, 1]$ that has the same distribution for all $b \in \mathbb{R}$ (this follows, for instance, from Theorem 3.1 of van der Vaart (1998)). It is well known that the best 5% level unbiased test in the Gaussian shift experiment is $\varphi_X^* = \mathbf{1}[|X/\sqrt{1 - \rho_0}| > \Phi(0.975)]$, so that the best unbiased 5% level test of H_0^s against $H_1^s(b)$ is $\varphi_{\rho_0}(G)$. Since $\varphi_{\rho_0}(G)$ does not depend on $\delta_0, \tilde{\beta}_0, \tilde{\delta}_0$, and $\rho_0, \delta_0, \tilde{\beta}_0, \tilde{\delta}_0$ were arbitrary, this proves the claim.

(ii) The bound $\bar{\pi}$ in Proposition 1 is constructed via Lemma 1, i.e., $\bar{\pi}$ is the power of the Neyman-Pearson test φ_{Λ^*} of H_{Λ^*} against H_F , treating G as the observation, with Λ^* the approximately least favorable distribution Λ^* determined as described in Section 3.1.

Now let $\tilde{\Lambda}^*$ be the probability distribution for $(\theta, \tilde{\beta}, \tilde{\delta})$ with the same marginal distribution of θ as Λ^* , and $(\tilde{\beta}, \tilde{\delta}) = (\beta A_{21}, \delta A_{21})$ conditional on θ . The Neyman-Pearson test of $H_{\tilde{\Lambda}^*}$ against $H_{\tilde{F}}$, based

on the observation $(G, \tilde{G}')'$, is then seen to be identical to φ_{Λ^*} . The result thus follows from Lemma 1.

(iii) The maximal invariant is $(\Psi, G(1))$, where $\Psi(s) = (G(s) - G(1), \tilde{G}(s)' - s\tilde{G}(1)')'$. Note that the distribution of $(\Psi, G(1))$ does not depend on $\tilde{\beta}$, and that $G(1)$ is independent of Ψ . The log of the RN derivative of $(\Psi, G(1))$ with respect to the distribution of (B_{k+1}, Z) , where B_{k+1} is a $k+1$ Brownian Bridge with covariance matrix Σ and $Z \sim \mathcal{N}(0, 1)$ independent of B_{k+1} , is equal to

$$\delta e_1' \Sigma^{-1} (\Psi(\rho) - \rho \Psi(1)) + (\beta + \delta \rho) G(1) - \frac{1}{2} \rho (1 - \rho) e_1' \Sigma^{-1} e_1 \delta^2 - \frac{1}{2} (\beta + \delta \rho)^2$$

where e_1 is the first column of I_{k+1} . By the same arguments as employed in part (i), this experiment is seen to be equivalent to the observation of $(\bar{\Psi}, G(1))$, where $\bar{\Psi}(s) = W(s) - sW(1) + (\rho(1-s) + \mathbf{1}[s \leq \rho])(s-\rho)\delta/\sqrt{1-R^2}$. A calculation yields that the RN derivative of the distribution of $(\bar{\Psi}, G(1))$ under $\theta \sim F$ with respect to the distribution of $(B_1, W(1))$ (where $B_1(s) = W(s) - sW(1)$) is

$$\frac{1}{0.70} \int_{0.15}^{0.85} \left(\frac{(1 + (1-r)\sigma_\beta^2)(1 + r(\sigma_\delta^2 - \sigma_\beta^2)) - R^2(1 + \sigma_\beta^2(1-2r) + r^2\sigma_\delta^2)}{1 - R^2} \right)^{-1/2} \\ \times \exp \left[\frac{1}{2} v(r)' \begin{pmatrix} 1 + \frac{1}{\sigma_\beta^2} - \frac{r(2-r)rR^2}{1-R^2} & -\frac{(1-r)rR^2}{1-R^2} \\ -\frac{(1-r)rR^2}{1-R^2} & \frac{1}{\sigma_\delta^2 - \sigma_\beta^2} + \frac{r(1-r)R^2}{1-R^2} \end{pmatrix}^{-1} v(r) \right] dr$$

where $v(r) = ((1-r)G(1) - \bar{\Psi}(r)/\sqrt{1-R^2}, rG(1) + \bar{\Psi}(r)/\sqrt{1-R^2})'$, and under $\delta \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $\rho = r$, it is

$$\left(1 + \frac{r(1-R^2r)\sigma_{\delta,1}^2}{1-R^2} \right)^{-1/2} \exp \left[-\frac{1}{2} \frac{\mu_1^2}{\sigma_{\delta,1}^2} + \frac{1}{2} \frac{\sigma_\delta^2 \left(\frac{\bar{\Psi}(\rho)}{\sqrt{1-R^2}} + rG(1) + \mu_1/\sigma_{\delta,1}^2 \right)^2}{1 + \sigma_{\delta,1}^2 \frac{r(1-R^2r)}{1-R^2}} \right].$$

The power bounds $\bar{\pi}_{inv}$ were computed via Lemma 1 using these densities as described in Section 3.4. The inequality $\bar{\pi}_{inv}(R^2) \leq \bar{\pi}_{inv}(\bar{R}^2)$ for $R^2 \leq \bar{R}^2$ holds because one can always reduce the information of the experiment with \bar{R}^2 to that with R^2 by adding a data independent Brownian Bridge of appropriate variance to $\bar{\Psi}$.

Condition 2 *In the stable model with parameter Γ_0*

(i) *in some neighborhood \mathcal{B}_0 of Γ_0 , $l_{T,t}(\Gamma)$ is twice differentiable a.s. with respect to Γ for $t = 1, \dots, T, T \geq 1$;*

(ii) *for all $s \in [0.15, 0.85]$ and $\epsilon > 0$ there exists $K(s, \epsilon) > 0$ such that $P(\sup_{\|\Gamma - \Gamma_0\| \geq \epsilon} T^{-1} \sum_{t=1}^{\lfloor sT \rfloor} (l_{T,t}(\Gamma) - l_{T,t}(\Gamma_0)) < -K(s, \epsilon)) \rightarrow 1$ and $P(\sup_{\|\Gamma - \Gamma_0\| \geq \epsilon} T^{-1} \sum_{t=\lfloor sT \rfloor + 1}^T (l_{T,t}(\Gamma) - l_{T,t}(\Gamma_0)) < -K(s, \epsilon)) \rightarrow 1$;*

(iii) *$\{s_{T,t}(\Gamma_0), \mathfrak{F}_{T,t}\}$ is a square-integrable martingale difference array with $T^{-1} \sum_{t=1}^{\lfloor sT \rfloor} E[s_{T,t}(\Gamma_0) s_{T,t}(\Gamma_0)' | \mathfrak{F}_{T,t-1}] \xrightarrow{p} s \Sigma^{-1}$ for all $0 \leq s \leq 1$ and some $(k+1) \times (k+1)$ dimensional matrix Σ , $T^{-1} \sup_{t \leq T} \|E[s_{T,t}(\Gamma_0) s_{T,t}(\Gamma_0)' | \mathfrak{F}_{T,t-1}]\| \xrightarrow{p} 0$ and there exists $\nu > 0$ such that $T^{-1} \sum_{t=1}^T E[\|s_{T,t}(\Gamma_0)\|^{2+\nu} | \mathfrak{F}_{T,t-1}] = O_p(1)$;*

- (iv) $T^{-1} \sum_{t=1}^T \|h_{T,t}(\Gamma_0)\| = O_p(1)$, $T^{-1} \sup_{t \leq T} \|h_{T,t}(\Gamma_0)\| \xrightarrow{p} 0$ and for any decreasing neighborhood \mathcal{B}_T of Γ_0 contained in \mathcal{B}_0 , $T^{-1} \sum_{t=1}^T \sup_{\Gamma \in \mathcal{B}_T} \|h_{T,t}(\Gamma) - h_{T,t}(\Gamma_0)\| \xrightarrow{p} 0$;
- (v) For all $0 \leq s \leq 1$, $T^{-1} \sum_{t=1}^{\lfloor sT \rfloor} h_{T,t}(\Gamma_0) \xrightarrow{p} -s\Sigma^{-1}$.

Part (ii) of Condition 2 is an identification condition that ensures consistency of the pre and post potential break maximum likelihood estimator—cf., for instance, Condition 6 on page 436 of Schervish (1995). Parts (ii)-(v) are a special case of Condition 2 in Li and Müller (2007) and Condition 1 in Müller and Petalas (2007), who provide further discussion and references.

Proof of Proposition 3:

We first show convergence of the experiment involving observation X_T as $T \rightarrow \infty$ to that of observing $(G, \tilde{G})'$ in the sense of Definition 9.1 of van der Vaart (1998).

By Lemma 1 of Li and Müller (2007), any unstable model with parameter evolution (16) is contiguous to the stable model. Furthermore, by the same reasoning as employed in the proof of Lemma 1 of Li and Müller (2007), under the stable model,

$$\ln \text{LR}_T = \int_0^1 \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix}' \Sigma^{-1} d\hat{W}_T(\lambda) - \frac{1}{2} \int_0^1 \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix}' \Sigma^{-1} \begin{pmatrix} \beta + \delta \mathbf{1}[\lambda \leq \rho] \\ \tilde{\beta} + \tilde{\delta} \mathbf{1}[\lambda \leq \rho] \end{pmatrix} d\lambda + o_p(1)$$

where $\hat{W}_T(\lambda) = T^{-1/2} \Sigma^{-1/2} \sum_{t=1}^{\lfloor \lambda T \rfloor} s_{T,t}(\Gamma_0)$. By Theorem 3.6 in McLeish (1974), under Condition 2 (iii), $\hat{W}_T(\cdot) \Rightarrow (W(\cdot), \tilde{W}_{k+1}(\cdot))$. Thus, by the Continuous Mapping Theorem (CMT), also $(\hat{W}_T(\cdot), \ln \text{LR}_T) \Rightarrow ((W(\cdot), \tilde{W}_{k+1}(\cdot)), \ln \text{LR})$ in the stable model, where $\ln \text{LR}$ is the r.h.s. of (18) with $(G, \tilde{G})'$ replaced by $(W, \tilde{W}'_{k+1})'$. Since LR is the RN derivative of the distribution of $(G, \tilde{G})'$ with respect to the distribution of $(W, (A_{21}W + A_{22}\tilde{W}_k)')$, the general version of LeCam's Third Lemma (see, for instance, Lemma 27 of Pollard (2001)) thus implies that $\hat{W}_T(\cdot) \Rightarrow (G(\cdot), \tilde{G}(\cdot))'$ in the unstable model (16). The convergence of the experiments now follows from the same arguments as employed in the proof of Theorem 9.4 of van der Vaart (1998).

Now for part (i), fix arbitrary $\theta_1 \in \Theta_1, \tilde{\beta}_1, \tilde{\delta}_1 \in \mathbb{R}^k$. Let T' be a subsequence of T such that $\limsup_{T \rightarrow \infty} E_{\theta_1, \tilde{\beta}_1, \tilde{\delta}_1} [\varphi_T(X_T)] = \lim_{T' \rightarrow \infty} E_{\theta_1, \tilde{\beta}_1, \tilde{\delta}_1} [\varphi_{T'}(X_{T'})]$. By (19), there is a further subsequence T'' and test φ such that $E_{\theta, \tilde{\beta}, \tilde{\delta}} [\varphi_{T''}(X_{T''})] \rightarrow E_{\theta, \tilde{\beta}, \tilde{\delta}} [\varphi(G, \tilde{G})]$ for all $\theta \in \Theta_0 \cup \Theta_1, \tilde{\beta}, \tilde{\delta} \in \mathbb{R}^k$. Since φ_T is asymptotically unbiased and of asymptotic level 5%, this implies φ to be a 5% level unbiased test. By Proposition 2(i), the most powerful unbiased test is φ_ρ , so that $\limsup_{T \rightarrow \infty} E_{\theta_1, \tilde{\beta}_1, \tilde{\delta}_1} [\varphi_T(X_T)] = E_{\theta_1, \tilde{\beta}_1, \tilde{\delta}_1} [\varphi(G, \tilde{G})] \leq E_{\theta_1, \tilde{\beta}_1, \tilde{\delta}_1} [\varphi_\rho(G)]$.

Parts (ii) and (iii) follow analogously.

Details on Determination of Approximate Least Favorable Distribution:

The distributions Ψ on Θ_0 mentioned in the main text are of the form " ρ is uniform on $[\max(0.15, a/100 - \frac{1}{2}), \min(0.85, b/100 + \frac{1}{2})]$ and δ is an equal probability mixture of $\delta \sim \mathcal{N}(\mu_\delta, \sigma_\delta^2)$ and $\delta \sim \mathcal{N}(-\mu_\delta, \sigma_\delta^2)$ " for some integers $a, b \in \{15, \dots, 85\}$, $b \geq a$, and

$\mu_\delta, \sigma_\delta^2 > 0$. Collect the four determining numbers of $\Psi = \Psi_\zeta$ in the vector $\zeta = (a, b, \mu_\delta, \sigma_\delta^2)$. The density of $G(\cdot)$ under Ψ_ζ , with respect to the distribution of $W(\cdot)$, is given by $\frac{1}{\min(0.85, b/100+1/2) - \max(0.15, a/100-1/2)} \int_{\max(0.15, a/100-1/2)}^{\min(0.85, b/100+1/2)} f(r, \mu_\delta, \sigma_\delta^2; G) dr$, where

$$f(r, \mu_\delta, \sigma_\delta^2; G) = (r\sigma_\delta^2 + 1)^{-1/2} \exp\left[\frac{1}{2} \frac{\sigma_\delta^2 G(r)^2 - r\mu_\delta^2}{r\sigma_\delta^2 + 1}\right] \cosh\left[\frac{G(r)^2 \mu_\delta}{r\sigma_\delta^2 + 1}\right]$$

and the density of $G(\cdot)$ with $(\beta, \delta, \rho) \sim F$ as in Condition 1 is given by $\frac{1}{0.70} \int_{0.15}^{0.85} h(r; G) dr$, where

$$h(r; G) = (r\sigma_\delta^2 + 1)^{-1/2} ((1-r)\sigma_\beta^2 + 1)^{-1/2} \exp\left[\frac{1}{2} \frac{\sigma_\delta^2 G(r)^2}{r\sigma_\delta^2 + 1} + \frac{1}{2} \frac{\sigma_\beta^2 (G(1) - G(r))^2}{(1-r)\sigma_\beta^2 + 1}\right].$$

As discussed in the main text, we approximate integrals with sums over the 100 point grid $r \in \{\frac{1}{100}, \frac{2}{100}, \dots, 1\}$. Thus, we replace $\frac{1}{0.70} \int_{0.15}^{0.85} h(r; G) dr$ by $\frac{1}{71} \sum_{l=15}^{85} h(l/100)$, and for the distribution Ψ_ζ , we approximate the resulting density by $f(\zeta; G) = \frac{1}{b-a+1} \sum_{l=a}^b f(l/100, \mu_\delta, \sigma_\delta^2; G)$ for $\zeta = (a, b, \mu_\delta, \sigma_\delta^2)$. All Λ we consider a mixtures of Ψ_ζ 's. Let $\zeta = (\zeta_1, \dots, \zeta_N)$, $\mathbf{p} = (p_1, \dots, p_N)$ with $\sum_{j=1}^N p_j = 1$. Thus, Neyman-Pearson tests can be written as $\varphi(\zeta, \mathbf{p}, cv) = \mathbf{1}[h(G)/\sum_{j=1}^N p_j f(\zeta_j; G) < 1/cv]$ for some $cv \in \mathbb{R}$, with null rejection probability $\pi_0(\zeta, \mathbf{p}, cv; \zeta) = \int E_\theta[\varphi(\zeta, \mathbf{p}, cv; G)] d\Psi_\zeta(\theta)$ under H_{Ψ_ζ} . In the following, write H_ζ for H_{Ψ_ζ} to ease notation. We approximate π_0 with the Monte Carlo estimator

$$\hat{\pi}_0(\zeta, \mathbf{p}, cv; \zeta) = \frac{1}{20000} \sum_{i=1}^{20000} \left(1 + \left(cv \sum_{j=1}^N p_j \frac{f(\zeta_j; G_i^*)}{h(G_i^*)} \right)^{10} \right)^{-1} \quad (35)$$

where G_i^* are (pseudo) random draws from G under Ψ_ζ . Because h and f only depend on $\{G(l/100)\}_{l=1}^{100}$, one can generate G_i^* by suitably transforming Gaussian Random Walks with 100 steps. In contrast to the standard Monte Carlo estimator based on averaging $\varphi(\zeta, \mathbf{p}, cv)$ directly, the numerically close analogue (35) is a differentiable function of (\mathbf{p}, cv) , which greatly facilitates the computations below.

The algorithm calls three subroutines SR1, SR2 and SR3.

SR1 The routine takes a set $\zeta = (\zeta_1, \dots, \zeta_N)$ and a levels $\alpha = (\alpha_1, \dots, \alpha_N)$ as given, and returns an estimate of the least favorable distribution Λ_ζ^* and corresponding critical value cv that describes the Neyman-Pearson test of the composite null hypothesis $H_\zeta : G$ has density Ψ_ζ for some $\zeta \in \zeta$ against H_F , which is of level α_j under H_{ζ_j} , $j = 1, \dots, N$. The distribution Λ_ζ^* is a mixture of the Ψ_ζ 's, $\zeta \in \zeta$. By the same argument as in Theorem 3.8.1 of Lehmann and Romano (2005), the least favorable mixing weights $\mathbf{p}^* = (p_1^*, \dots, p_N^*)$ have the two properties (i) $\pi_0(\zeta, \mathbf{p}^*, cv; \zeta) \leq \alpha_l$ for $l = 1, \dots, N$; and (ii) $\pi_0(\zeta, \mathbf{p}^*, cv; \zeta_l) < \alpha$ only if $p_l = 0$. This motivates the joint determination of \mathbf{p} and cv as numerical solutions to

$$\hat{\pi}_0(\zeta, \mathbf{p}, cv; \zeta_l) \leq \alpha_l \quad \text{and} \quad p_l(\hat{\pi}_0(\zeta, \mathbf{p}, cv; \zeta_l) - \alpha_l) = 0 \quad \text{for } l = 1, \dots, N. \quad (36)$$

Specifically, we determine appropriate \mathbf{p} and cv by minimizing the objective function

$$\sum_{l=1}^N (a_0 p_l + \exp[a_1(\hat{\pi}_0(\zeta, \mathbf{p}, cv; \zeta_l) - \alpha_l)])(\hat{\pi}_0(\zeta, \mathbf{p}, cv; \zeta_l) - \alpha_l)^2 \quad (37)$$

where $a_0 = 100$ and $a_1 = 2000$. As a function of \mathbf{p} and cv , (37) is continuous and with known first derivative, so that a standard quasi-Newton optimizer can be employed. Also, the $20000N^2$ numbers $f(\varsigma_l; G_i^*)/h(G_i^*)$ for G_i^* drawn under ς_j for $i = 1, \dots, N$, $l = 1, \dots, N$ and $j = 1, \dots, m$ can be computed and stored once to speed up the the evaluation of $\hat{\pi}_0(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv}; \varsigma_l)$ and its partial derivatives.

SR2 The routine takes $(\boldsymbol{\varsigma}, \mathbf{p}, \boldsymbol{\alpha})$ as inputs and returns $(\boldsymbol{\varsigma}', \mathbf{p}', \boldsymbol{\alpha}')$ of length $N' \leq N$ by eliminating values $(\theta_j, p_j, \alpha_j)$ with p_j approximately equal to zero.

SR3 The routine takes a test φ described by the triple $(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$, a set grid of values $\boldsymbol{\varsigma}_g = (\varsigma_{g,1}, \dots, \varsigma_{g,N_g})$ and a nominal level $\alpha_n \in [0, 1]$ as given, and determines whether $\varphi(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$ is of level α_n under $H_{\varsigma'}$, $\varsigma' \in \boldsymbol{\varsigma}'$. Specifically, it evaluates $\hat{\pi}_0(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv}; \varsigma')$ for all $\varsigma' \in \boldsymbol{\varsigma}'$ in a random order until $\hat{\pi}_0(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv}; \varsigma') > \alpha_n$, in which case ς' is returned. Otherwise, the routine returns a flag indicating that the test is of level α_n .

Overall the algorithm iterates between the subroutines as follows:

1. Initialize $\boldsymbol{\varsigma} = ((15, 24, 28, 200), (75, 85, 28, 200), (15, 85, 20, 4), (85, 85, 3, 4), (15, 85, 0, 300))$ and $\boldsymbol{\alpha} = (0.033, 0.028, 0.034, 0.0445)$. Call SR1 and SR2 to obtain $(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$.
2. For $i = 1$ to 7:
 - (a) Call SR3 with $(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$, $\alpha_n = \alpha_{n,i}$ and $\boldsymbol{\varsigma}_g = \boldsymbol{\varsigma}_{g,i}$.
 - (b) If SR3 returns ς , extend $\boldsymbol{\varsigma}$ by ς and $\boldsymbol{\alpha}$ by α_i , and call SR1 and SR2 to obtain new $(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$. Go to step a.
 - (c) If SR3 returns flag that $\varphi(\boldsymbol{\varsigma}, \mathbf{p}, \text{cv})$ is of level α_n , increase i by one.

The 7 grids $\boldsymbol{\varsigma}_{g,i}$ and $\alpha_{n,i}$ are described in Table 5. The grid contains all possible combinations of $\varsigma = (a, b, \mu_\delta, \sigma_\delta^2)$ with a and b consecutive values in column 2, and μ_δ as in column 3. The randomization in SR3 prevents cycles that arise through the thinning operation in SR2.

Details on Size Control Computations:

The test φ^* as defined above only depends on $\{G(l/100)\}_{l=1}^{100}$, and conditional on the Monte Carlo draw of the random walk $\{W_i(l)\}_{l=1}^{100}$, $G_i(l/100) = W_i(l/100) + \delta \min(\rho, l/100)$ is a function of δ and ρ . Write $S_i(\delta, \rho)$ for the induced function $\varphi^*(G_i)$. The approach outlined in Section 3.3 requires that for any given values $0 \leq \delta_1 < \delta_2$ and $0.15 \leq \rho_1 < \rho_2 \leq 0.85$, an algorithm must (attempt to) determine $\min_{\delta_1 \leq \delta < \delta_2, \rho_1 \leq \rho < \rho_2} S_i(\delta, \rho)$ and $\max_{\delta_1 \leq \delta < \delta_2, \rho_1 \leq \rho < \rho_2} S_i(\delta, \rho)$ (it is sufficient to consider non-negative δ , because $\varphi^*(G) = \varphi^*(-G)$). For this determination, we now consider how supF_0 , \hat{l}_0 , \hat{t}_0 and LR_0 behave as functions of (δ, ρ) on the rectangles $\delta_1 \leq \delta < \delta_2, \rho_1 \leq \rho < \rho_2$ where ρ_1 and ρ_2 are such that $l_0 - 1 \leq 100\rho_1$ and $100\rho_2 < l_0$ for some $l_0 \in \{16, \dots, 85\}$, conditional on W_i .

Table 5: Grids for Computation of Approximate Least Favorable Distribution

| i | l | μ_δ | σ_δ^2 | α_i |
|-----|--|---------------------------------|-------------------|------------|
| 1 | 15, 20, 75, 83, 85 | 0, 2, \dots , 20 | 300 | 4.45% |
| 2 | 15, 20, 75, 83, 85 | 0, 2, \dots , 16 | 50 | 4.45% |
| 3 | 15, 20, 75, 83, 85 | 0, 2, \dots , 22 | 10 | 4.50% |
| 4 | 15, 20, 75, 83, 85 | 0, 2, \dots , 22 | 3 | 4.55% |
| 5 | 15, 20, 30, 45, 60, 70, 75, 80, 83, 85 | 0, 2, \dots , 22 | 10 | 4.55% |
| 6 | 83, 84, 85 | 0, $\frac{1}{2}$, \dots , 18 | 1 | 4.60% |
| 7 | 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 83, 85 | 0, 1, \dots , 22 | 10 | 4.60% |

For supF_0 , note that $((100-l)G(\frac{l-1}{100}) - (l-1)(G(1) - G(\frac{l}{100}))) / \sqrt{(100-l)(l-1)} = a^{\text{supF}}(l, W_i) + \delta \cdot b^{\text{supF}}(l, \rho)$, where $a^{\text{supF}}(l, W_i)$ does not depend on (ρ, δ) , and $b^{\text{supF}}(l, \rho) = b_{pre}^{\text{supF}}(l, \rho) = \sqrt{(l-1)/(100-l)}(1-\rho)$ for $l < l_0$, $b^{\text{supF}}(l, \rho) = b_{mid}^{\text{supF}}(l) = \sqrt{(100-l)(l-1)}/100$ for $l = l_0$ and $b^{\text{supF}}(l, \rho) = b_{post}^{\text{supF}}(l, \rho) = \sqrt{(100-l)/(l-1)}\rho$ for $l > l_0$. Thus, as long as $\frac{100}{99}a(l, W_i)^2 < 90$ (which is true for almost all draws of W_i),

$$\begin{aligned} \delta_1 \left(\max_{16 \leq l < l_0} b_{pre}(l, \rho_2) + b_{mid}(l_0) + \max_{l_0 < l \leq 85} b_{post}(l, \rho_1) \right) &\leq \sqrt{\frac{100}{99} \text{supF}_0} \\ &\leq \delta_2 \left(\max_{16 \leq l < l_0} b_{pre}(l, \rho_1) + b_{mid}(l_0) + \max_{l_0 < l \leq 85} b_{post}(l, \rho_2) \right). \end{aligned}$$

For \hat{l}_0 , note that with $\xi = \rho\delta$, $S_0(l)$ is of the form $S_0(l) = S_0(l, \delta, \xi) = \sum_{j=1}^2 (a_j^S(l, W_i) + b_{i,j}^S(l)\delta + c_{i,j}^S(l)\xi)^2$ for $i \in \{pre, mid, post\}$ in self-explanatory notation. In particular, $b_{pre,1}^S(l) = l$, $b_{post,1}^S(l) = b_{mid,1}^S(l) = 0$, $b_{pre,2}^S(l) = 0$, $b_{mid,2}^S(l) = (l-1)l/100$, $b_{post,2}^S(l) = 0$, $c_{pre,1}^S(l) = -l$, $c_{post,1}^S(l) = c_{mid,1}^S(l) = 100-l$, $c_{pre,2}^S(l) = 0$, $c_{mid,2}^S(l) = -(l-1)$, and $c_{post,2}^S(l) = 1$. Let $\bar{\rho} = \frac{1}{2}(\rho_1 + \rho_2)$, $\bar{\delta} = \frac{1}{2}(\delta_1 + \delta_2)$, and $\bar{\xi} = \bar{\rho}\bar{\delta}$, and define $\hat{l} = \arg \max_{16 \leq l \leq 85} S_0(l, \bar{\delta}, \bar{\xi})$. With $S_\Delta(l, \delta, \xi) = S_0(l, \delta, \xi) - S_0(\hat{l}, \bar{\delta}, \bar{\xi})$ and $\mathcal{T} = [\delta_1, \delta_2] \times [\xi_1, \xi_2]$, $\xi_1 = \delta_1\rho_1$, $\xi_2 = \delta_2\rho_2$, the set of possible values for \hat{l}_0 is a (possibly proper) subset of $\hat{L}_0 = \{16 \leq l \leq 85 : \sup_{(\delta, \xi) \in \mathcal{T}} S_\Delta(l, \delta, \xi) \geq 0\}$, because the range of $S_\Delta(l, \delta, \xi)$ on \mathcal{T} is at least as large as the range of $S_\Delta(l, \delta, \rho\delta)$ on $[\delta_1, \delta_2] \times [\rho_1, \rho_2]$, for each l . Since $S_\Delta(l, \delta, \xi)$, viewed as a function of δ and ξ , is a quadratic polynomial, the minimum and maximum occur either in a corner $(\delta, \xi) \in \{(\delta_1, \xi_1), (\delta_1, \xi_2), (\delta_2, \xi_1), (\delta_2, \xi_2)\}$, or, possibly, at a local maximum along one of the four sides, or, possibly, at an interior extremum. The five potential interior local extrema are easily computed by solving the appropriate linear first order conditions.

Given \hat{L}_0 , it is straightforward to construct upper and lower bounds on \hat{t}_0 .

Finally, for LR_0 , note that the exponents in the numerator is a positive definite quadratic polynomial in δ and $\xi = \rho\delta$. Expanding $\cosh(x) = \frac{1}{2}\exp(x) + \frac{1}{2}\exp(-x)$, also the denominator can be written as a sum of exponentials where each exponent is a positive definite quadratic polynomial in (δ, ξ) . Since positive definite quadratic polynomials are convex, and the sum of log-convex functions is log convex, both the numerator and denominator are log-convex functions of (δ, ξ) , say, $N_{\text{LR}}(\delta, \xi)$

and $D_{\text{LR}}(\delta, \xi)$. Let \bar{v}_D be the 2×1 vector of partial derivatives of $\ln D_{\text{LR}}(\delta, \xi)$ at $(\delta, \xi) = (\bar{\delta}, \bar{\xi})$. Since $\ln D_{\text{LR}}(\delta, \xi) - (\delta - \bar{\delta}, \xi - \bar{\xi})\bar{v}_D$ is convex with zero derivative at $(\delta, \xi) = (\bar{\delta}, \bar{\xi})$, it takes on its minimum at $(\bar{\delta}, \bar{\xi})$, and since $\ln N_{\text{LR}}(\delta, \xi) - (\delta - \bar{\delta}, \xi - \bar{\xi})\bar{v}_D$ is convex, it takes on its maximum in one of the corners. Thus,

$$\max_{\delta_1 \leq \delta < \delta_2, \rho_1 \leq \rho < \rho_2} \text{LR}_0 \leq \frac{\max_{(\delta, \xi) \in \mathcal{T}} N_{\text{LR}}(\delta, \xi) \exp[-(\delta - \bar{\delta}, \xi - \bar{\xi})\bar{v}_D]}{D_{\text{LR}}(\bar{\delta}, \bar{\xi})}$$

and, reversing the role of the numerator and denominator, we can construct an analogous lower bound.

Details on Power Bound Computations:

A standard Wiener process on the unit interval W can be decomposed as $W(s) = D_W^\tau(s) + \alpha_W + (s - 1/2)\beta_W$, where α_W and β_W are the coefficients of a continuous time least squares regression of $W(s)$ on $(1, s - 1/2)$, and $D_W^\tau(s)$ are the residuals. Let $D_W^{\text{sq}} = \int D^\tau(s)^2 ds$. With these definitions, a straightforward calculation yields $\int_0^1 W(s)^2 ds = D_W^{\text{sq}} + \frac{1}{3}\beta_W^2 + \alpha_W\beta_W + \alpha_W^2$. Further, define $D_W(s)$ as the demeaned Brownian Bridge $W(s) - sW(1)$, $D_W(s) = W(s) - sW(1) - \alpha_W + \frac{1}{2}W(1)$, and note that $\Delta_W = \sup_{0 \leq s \leq 1} |W(s) - \alpha_W| \leq \frac{1}{2}|W(1)| + \sup_{0 \leq s \leq 1} |D_W(s)|$ and $\Delta_W^{\text{sq}} = \sup_{0 \leq s \leq 1} W(s)^2 \leq 2\Delta_W|\alpha_W| + \Delta_W^2 + |\int_0^1 W(s)^2 ds - \alpha_W^2|$. These expressions are useful because (i) $(W(1), \alpha_W, \beta_W)$ are jointly Gaussian with known covariance matrix and independent of $D_W^\tau(\cdot)$ (and thus D_W^{sq}); (ii) the distribution of D_W^{sq} can be computed exactly by inverting the characteristic function derived in Theorem 9.12 of Tanaka (1996); (iii) Darling (1983) provides a formula for the c.d.f. of $\sup_{0 \leq s \leq 1} D_W(s)$.

Evaluation of the expression in Darling (1983) yields $P(\sup_{0 \leq s \leq 1} D(s) > 2.2) < \varepsilon_D = 5 \cdot 10^{-12}$, so that $P(\sup_{0 \leq s \leq 1} |D_W(s)| > 2.2) < 2\varepsilon_D$. Also, evaluation of the c.d.f. of D_W^{sq} at the 15 points $0.1, 0.2, \dots, 1.5$ by inverting the characteristic function, as discussed in Chapter 6 of Tanaka (1996), permits the straightforward construction of the two random variables $D_W^{\text{sq}^l}$ and $D_W^{\text{sq}^u}$ such that $P(D_W^{\text{sq}^l} \leq D_W^{\text{sq}} \leq D_W^{\text{sq}^u}) = 1$ by suitably transforming uniform random variable on the unit interval (with $D_W^{\text{sq}^u}$ taking on the value $+\infty$ with probability $P(D_W^{\text{sq}} > 1.5) \simeq 4 \cdot 10^{-14}$). Let $(Z_j, \alpha_{W_j}, \beta_{W_j})$ be J i.i.d. Monte Carlo draws from the distribution of $(W(1), \alpha_W, \beta_W)$, let $(D_j^{\text{sq}^l}, D_j^{\text{sq}^u})$ be J independent i.i.d. Monte Carlo draws from the distribution of $(D^{\text{sq}^l}, D^{\text{sq}^u})$. Then define $S_{1,j}^u = 2.2 + 0.5|Z_j|$ and $S_{2,j}^u = 2|\alpha_{W_j}|S_{1,j}^u + (S_{1,j}^u)^2 + |D_{W_j}^{\text{sq}^u} + \frac{1}{3}\beta_{W_j}^2 + \alpha_{W_j}\beta_{W_j}|$, $W_{b,j+1} = W_{b,j} + Z_j/\sqrt{J}$, $\bar{W}_j = W_{b,j} + \alpha_W/\sqrt{J}$, $\bar{W}_j^{2^l} = (D_{W_j}^{\text{sq}^l} + \frac{1}{3}\beta_{W_j}^2 + \alpha_{W_j}\beta_{W_j})/J + \alpha_{W_j}W_{b,j}/\sqrt{J} + W_{b,j}^2$ and $\bar{W}_j^{2^u} = \bar{W}_j^{2^l} + (D_{W_j}^{\text{sq}^u} - D_{W_j}^{\text{sq}^l})/J$ for $j = 1, \dots, J$, where $W_{b,1} = 0$. With the implemented number of $J = 800,000$ intervals, a draw $\{\bar{W}_j^{2^l}, \bar{W}_j^{2^u}, \bar{W}_j, S_{1,j}^u, S_{2,j}^u\}_{j=1}^J$ satisfies $\bar{W}_j^{2^l} \leq \bar{W}_j^2 \leq \bar{W}_j^{2^u}$, $S_{1,j} \leq S_{1,j}^u$ and $S_{2,j} \leq S_{2,j}^u$ for all $j = 1, \dots, J$ with probability of at least $(1 - 2\varepsilon_D)^J > 1 - 7 \cdot 10^{-5}$ by construction. Thus, very slight corrections to the estimated critical value and Monte Carlo rejection probability yields an unconditionally conservative Monte Carlo estimator of the power bounds.

Proof of Proposition 4:

(i) The CMT, and, in the case of \hat{l} , some tedious algebra, imply that $\hat{l} \Rightarrow \hat{l}_0$ and $\widehat{\text{supF}} \Rightarrow \text{supF}_0$ with \hat{l}_0 and supF_0 as defined the definition of φ^* above (note that replacing G by G_0 leaves \hat{l}_0 and supF_0

unchanged). Define $\beta_T = T^{1/2}(\gamma_{post} - \gamma_{post,0})$ and $\delta_T = T^{1/2}(\gamma_{pre} - \gamma_{post})$.

(i.a) Since $G(s) = G_0(s) + \beta s$, the result follows from the CMT.

(i.b) We have $\hat{t}_{post}/\beta_T \xrightarrow{P} \sqrt{1 - l_0/100}/\sigma$ and a calculation yields $\beta_T^{-2} \ln \widehat{\text{LR}} \xrightarrow{P} c > 0$, so that $P(|\hat{t}_{post}| > 2.01) \rightarrow 1$ and $P(\widehat{\text{LR}} > 2.41) \rightarrow 1$.

(ii) Note that $\delta_T^{-2} (\hat{\gamma}_{post}(\lceil 100\rho \rceil) - \hat{\gamma}_{pre}(\lceil 100\rho \rceil - 1))^2 \xrightarrow{P} 1$ and $\hat{\omega}_{post}^2(\lceil 100\rho \rceil) \xrightarrow{P} \sigma^2/(1 - l/100)$, so that $P(\widehat{\text{supF}} > 90) \rightarrow 1$. Let

$$\hat{S}(l) = \sum_{j=1}^{l-1} (\hat{\Delta}_{pre}(j) - \hat{\gamma}_{pre}(l-1))^2 + \sum_{j=l+1}^{100} (\hat{\Delta}_{post}(j) - \hat{\gamma}_{post}(l))^2$$

where $\hat{\Delta}_{pre}(j) = \hat{\gamma}_{pre}(15)$ for $j \leq 15$ and $\hat{\Delta}_{post}(j) = \hat{\gamma}_{post}(85)$ for $j > 85$, and note that $\hat{l} = \arg \max_{16 \leq l \leq 85} \hat{S}(l)$. If $100\rho \neq \lceil 100\rho \rceil$, then $\hat{S}(\lceil 100\rho \rceil + 1) = O_p(T^{-1})$, and for all $l \neq \lceil 100\rho \rceil + 1$, $P(T\hat{S}(l) > M) \rightarrow 1$ for any $M \in \mathbb{R}$, so that $\hat{l} \xrightarrow{P} \lceil 100\rho \rceil + 1$. Part (ii.b), and part (ii.a) for $\rho \notin R_{grid}$, now follow from the CMT.

Consider thus the proof of part (ii.a) if $100\rho = \lceil 100\rho \rceil$. By the same argument, $P(\hat{l} \notin \{l_0, l_0 + 1\}) \rightarrow 0$, where $l_0 = 100\rho$. If $l_0 \in \{84, 85\}$, the result follows again from the CMT. Otherwise, define $t_{post}(l_0 + 1) = (W(1) - W(\frac{l_0+1}{100}))/\sqrt{1 - \frac{l_0+1}{100}} \sim \mathcal{N}(0, 1)$ and $t_{post}(l_0 + 2) = (W(1) - W(\frac{l_0+2}{100}))/\sqrt{1 - \frac{l_0+2}{100}} \sim \mathcal{N}(0, 1)$, and consider

$$\begin{aligned} \hat{S}(l_0) - \hat{S}(l_0 + 1) &= A_{pre} + \hat{\Delta}_{post}(l_0 + 1)^2 - (100 - l_0)\hat{\gamma}_{post}(l_0)^2 + (99 - l_0)\hat{\gamma}_{post}(l_0 + 1)^2 \\ &\Rightarrow \Delta S(l_0) = a \cdot t_{post}(l_0 + 1)^2 + t_{post}(l_0 + 1)B(l_0) + C(l_0) \end{aligned}$$

where A_{pre} is a function of $\{\hat{\gamma}_{pre}(l)\}_{l=15}^{l_0-1}$, the convergence follows from the CMT, a is a positive constant, and $B(l_0)$ and $C(l_0)$ are random variables that are independent of $t_{post}(l_0 + 1)$ and $t_{post}(l_0 + 2)$. Thus, by a further application of the CMT, we obtain

$$\hat{t}_{post} \Rightarrow \mathbf{1}[\Delta S(l_0) \geq 0] \mathbf{1}[|t_{post}(l_0 + 1)| > 2.01] + \mathbf{1}[\Delta S(l_0) < 0] \mathbf{1}[|t_{post}(l_0 + 2)| > 2.01]. \quad (38)$$

Consider now the rejection probability of the r.h.s. of (38) conditional on $B(l_0) = b$ and $C(l_0) = c$. Noting that $\Delta S(l_0)$ is a quadratic function of $t_{post}(l_0 + 1)$ with positive coefficient on the square term, one obtains that the rejecting probability is bounded from above by the replacement $\mathbf{1}[\Delta S(l_0) \geq 0] = \mathbf{1}[t_{post}(l_0 + 1) < -2.01]$, for any value of b, c . A numerical calculation now shows that the conditional rejection probability remains below 5% even in that case for all $l_0 = 15, \dots, 83$, and the result follows.

(iib) Immediate from $P(\hat{l} \notin \{l_0, l_0 + 1\}) \rightarrow 0$ and $\beta_T^{-1}(\hat{\gamma}_{post}(l) - \gamma_{post,0})/\hat{\omega}_{post}(l) \xrightarrow{P} \sqrt{1 - l/100}/\sigma > 0$ for all $l \geq l_0$.

(iii) The claim is essentially a special case of Theorem 1 of Andrews and Guggenberger (2007c). Let θ_T be a parameter sequence such that $\limsup_{T \rightarrow \infty} E_{\theta_T}[\hat{\varphi}^*(Y_T)] = \limsup_{T \rightarrow \infty} \sup_{\theta \in \Theta_0} E_{\theta}[\hat{\varphi}^*(Y_T)]$. Pick a subsequence T' of T such that $\lim_{T' \rightarrow \infty} E_{\theta_{T'}}[\hat{\varphi}^*(Y_{T'})] = \limsup_{T \rightarrow \infty} E_{\theta_T}[\hat{\varphi}^*(Y_T)]$. Since $\bar{\Theta}_0 = \Theta_0 \cup \{(0, \rho, \delta) : \rho \in [0.15, 0.85], \delta \in \{+\infty, -\infty\}\}$ is compact under an appropriate metric, there exists a further subsequence $\theta_{T''}$ of $\theta_{T'}$ such that $\theta_{T''} \rightarrow \tilde{\theta} = (0, \tilde{\rho}, \tilde{\delta}) \in \bar{\Theta}_0$. If $\tilde{\rho} \notin R_{grid}$, then the result

follows as in parts (i) and (ii). If $\tilde{\rho} \in R_{grid}$, then by assumption, there is a further subsequence such that (32) holds, except for (33). Proceeding as in the proof of part (ii.a) now again yields the result (and for the case $\nu_{pre}(l_0) \in \{+\infty, -\infty\}$ or $\nu_{post}(l_0) \in \{+\infty, -\infty\}$, $P(|\Delta S(l_0)| < M) \rightarrow 0$ for all M , so that $\mathbf{1}[\Delta S(l_0) \geq 0] \xrightarrow{P} 0$ or $\mathbf{1}[\Delta S(l_0) \geq 0] \xrightarrow{P} 1$).

Justification of (31) via Andrews (1993):

The true parameters are given by $\Gamma_{T,t} = \Gamma_{post} + T^{-1/2} \mathbf{1}[T \leq 100\rho_T] \Delta_T$, where $\Delta_T = -(\omega\delta_T, -\tilde{\delta}_T) \rightarrow \Delta_0 \in \mathbb{R}^{k+1}$ and $\rho_T \rightarrow \rho_0 \in [0.15, 0.85]$, so that $E[g(X_{T,t}, \Gamma_{T,t})] = 0$. Write $g_t(\Gamma)$ for $g(X_{T,t}, \Gamma)$. Our set-up corresponds to what Andrews (1993) refers to as 'pure' structural change. We now show that under Assumption 1 of Andrews (1993), with the assumption in part (b) $E[g_t(\Gamma_0)] = 0$ replaced by $E[g_t(\Gamma_{T,t})] = 0$, (31) holds. These computations closely correspond to what is derived in Section 5.4 of Andrews (1993), with two differences: first, we do not assume that $\max_{t \leq T} \sup_{\|\Gamma - \Gamma_{post}\| \leq T^{-1/2}K} \|E[\Upsilon_t(\Gamma)] - \tilde{\Upsilon}\| \rightarrow 0$ for some $K > 0$ (cf. first line on page 832), and second, we consider sequences for Δ_T and ρ_T . (We note that there is a typo in Assumption 1-LP of Andrews (1993); it should read $\sup_{\pi \in \Pi} \|\sqrt{T}E\bar{m}_T(\theta_0, \pi) - \mu(\pi)\| \rightarrow 0$.)

By the argument in Andrews' proof of Theorem 4 (a), it suffices to show that $\sup_{0 \leq s \leq 1} \|T^{-1/2}E[\sum_{t=1}^{\lfloor sT \rfloor} g_t(\Gamma_{post}) + \tilde{\Upsilon} \int_0^s \eta_0(l)dl]\| \rightarrow 0$. Now clearly $\sup_{0 \leq s \leq 1} \|\lim_{T \rightarrow \infty} T^{-1} \sum_{t=1}^{\lfloor sT \rfloor} \eta_T(t/T) - \int_0^s \eta_0(l)dl\| \rightarrow 0$. Also,

$$\begin{aligned} T^{-1/2}E\left[\sum_{t=1}^{\lfloor sT \rfloor} g_t(\Gamma_{post}) + T^{-1/2}\tilde{\Upsilon}\eta_T(t/T)\right] &= T^{-1/2}E\left[\sum_{t=1}^{\lfloor \min(s, \rho_T)T \rfloor} g_t(\Gamma_{post}) + T^{-1/2}\tilde{\Upsilon}\eta_T(t/T)\right] \\ &= (\tilde{\Upsilon} - E[\tilde{\Upsilon}_T(\min(s, \rho_T))])\Delta_T \end{aligned}$$

where $\tilde{\Upsilon}_T(r)$ is equal to $T^{-1} \sum_{t=1}^{\lfloor rT \rfloor} \int_0^1 \Upsilon_t(\Gamma_{post} + \lambda T^{-1/2} \Delta_T) d\lambda$. Furthermore,

$$\begin{aligned} \sup_{0 \leq s \leq 1} \|E[\tilde{\Upsilon}_T(\min(s, \rho_T)) - \tilde{\Upsilon}_T(\min(s, \rho_0))]\| &\leq T^{-1}E \sum_{t=\lfloor \min(\rho_0, \rho_T)T \rfloor}^{\lfloor \max(\rho_0, \rho_T)T \rfloor} \left\| \int_0^1 \Upsilon_t(\Gamma_{post} + \lambda T^{-1/2} \Delta_T) d\lambda \right\| \\ &\leq (\lfloor \max(\rho_0, \rho_T)T \rfloor - \lfloor \min(\rho_0, \rho_T)T \rfloor) \sup_{t \leq T} E[\sup_{\Gamma \in \mathcal{G}_0} \|\Upsilon_t(\Gamma)\|] \rightarrow 0 \end{aligned}$$

where the convergence follows, since by Andrews' assumption 1(f), $\sup_{t \leq T} E[\sup_{\Gamma \in \mathcal{G}_0} \|\Upsilon_t(\Gamma)\|] = O(1)$ for some neighborhood \mathcal{G}_0 of Γ_{post} . Finally, $\sup_{0 \leq s \leq 1} \|\tilde{\Upsilon} - E[\tilde{\Upsilon}_T(\min(s, \rho_0))]\| \rightarrow 0$ follows from Andrews' (A.8).

References

- ANDERSON, R. J., AND K. L. KLIESON (2006): “The 1990’s Acceleration in Labor Productivity: Causes and Measurement,” *Federal Reserve Bank of St. Louis Review*, 88, 181–202.
- ANDREWS, D. W. K. (1991): “Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimation,” *Econometrica*, 59, 817–858.
- (1993): “Tests for Parameter Instability and Structural Change with Unknown Change Point,” *Econometrica*, 61, 821–856.
- ANDREWS, D. W. K., AND P. GUGGENBERGER (2007a): “Applications of Subsampling, Hybrid, and Size-Correction Methods,” *Cowles Foundation Discussion Paper No. 1608*.
- (2007b): “Hybrid and Size-Corrected Subsample Methods,” *Cowles Foundation Discussion Paper No. 1606*.
- (2007c): “The Limit of Finite-Sample Size and a Problem with Subsampling,” *Cowles Foundation Discussion Paper No. 1605*.
- ANDREWS, D. W. K., M. J. MOREIRA, AND J. H. STOCK (2006): “Optimal Two-Sided Invariant Similar Tests for Instrumental Variables Regression,” *Econometrica*, 74, 715–752.
- (2007): “Efficient Two-Sided Nonsimilar Invariant Tests in IV Regression with Weak Instruments,” *Working paper, Harvard University*.
- ANDREWS, D. W. K., AND W. PLOBERGER (1994): “Optimal Tests When a Nuisance Parameter Is Present Only under the Alternative,” *Econometrica*, 62, 1383–1414.
- BAI, J. (1994): “Least Squares Estimation of a Shift in Linear Processes,” *Journal of Time Series Analysis*, 15, 453–470.
- (1997): “Estimation of a Change Point in Multiple Regressions,” *Review of Economics and Statistics*, 79, 551–563.
- BAI, J., AND P. PERRON (1998): “Estimating and Testing Linear Models with Multiple Structural Changes,” *Econometrica*, 66, 47–78.
- BUNZEL, H., AND T. J. VOGELSANG (2005): “Powerful Trend Function Tests That are Robust to Strong Serial Correlation with an Application to the Prebisch-Singer Hypothesis,” *Journal of Business and Economic Statistics*, 23, 381–394.

- CHAMBERLAIN, G. (2000): “Econometric Applications of Maximin Expected Utility,” *Journal of Applied Econometrics*, 15, 625–644.
- COGLEY, T., AND T. J. SARGENT (2005): “Drifts and Volatilities: Monetary Policies and Outcomes in the Post WWII US,” *Review of Economic Dynamics*, 8, 262–302.
- DARLING, D. A. (1983): “On the Supremum of a Certain Gaussian Process,” *The Annals of Probability*, 11, 803–806.
- DUFOUR, J.-M. (2006): “Monte Carlo Tests with Nuisance Parameters: A General Approach to Finite-Sample Inference and Nonstandard Asymptotics,” *Journal of Econometrics*, 133, 443–477.
- ELLIOTT, G., AND U. K. MÜLLER (2006): “Efficient Tests for General Persistent Time Variation in Regression Coefficients,” *Review of Economic Studies*, 73, 907–940.
- (2007): “Confidence Sets for the Date of a Single Break in Linear Time Series Regressions,” *Journal of Econometrics*, 141, 1196–1218.
- GORDON, R. J. (2003): “Exploding Productivity Growth: Context, Causes, and Implications,” *Brookings Papers on Economic Activity*, 2, 207–279.
- GUGGENBERGER, P. (2008): “The Impact of a Hausman Pretest on the Size of Hypothesis Tests,” *Cowles Foundation Discussion Paper No. 1651*.
- HALL, A., S. HAN, AND O. BOLDEA (2008): “Inference Regarding Multiple Structural Changes in Linear Models Estimated Via Two Stage Least Squares,” *MPRA working paper 9251*.
- HANSEN, B. E. (2000): “Testing for Structural Change in Conditional Models,” *Journal of Econometrics*, 97, 93–115.
- JANSSON, M., AND M. J. MOREIRA (2006): “Optimal Inference in Regression Models with Nearly Integrated Regressors,” *Econometrica*, 74, 681–714.
- JORGENSON, D., M. S. HO, AND K. J. STIROH (2008): “A Retrospective Look at the US Productivity Growth Resurgence,” *Journal of Economic Perspectives*, 22, 3–24.
- KING, M. L. (1988): “Towards a Theory of Point Optimal Testing,” *Econometric Reviews*, 6, 169–218.

- LEHMANN, E. L., AND J. P. ROMANO (2005): *Testing Statistical Hypotheses*. Springer, New York.
- LI, H., AND U. K. MÜLLER (2007): “Valid Inference in Partially Unstable GMM Models,” *Working paper, Princeton University*.
- MCLEISH, D. L. (1974): “Dependent Central Limit Theorems and Invariance Principles,” *The Annals of Probability*, 2, 620–628.
- MÜLLER, U. K. (2007): “An Alternative Sense of Asymptotic Efficiency,” *Working paper, Princeton University*.
- MÜLLER, U. K., AND P. PETALAS (2007): “Efficient Estimation of the Parameter Path in Unstable Time Series Models,” *Working paper, Princeton University*.
- MÜLLER, U. K., AND M. W. WATSON (2008a): “Low-Frequency Robust Cointegration Testing,” *Working paper, Princeton University*.
- (2008b): “Testing Models of Low-Frequency Variability,” *Econometrica*, 76, 979–1016.
- NEWBY, W. K., AND K. WEST (1987): “A Simple, Positive Semi-Definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix,” *Econometrica*, 55, 703–708.
- ORPHANIDES, A. (2004): “Monetary Policy Rules, Macroeconomic Stability, and Inflation: A view from the Trenches,” *Journal of Money, Credit, and Banking*, 36, 151–175.
- POLLARD, D. (2001): “Contiguity,” Working paper, Yale University.
- SCHERVISH, M. J. (1995): *Theory of Statistics*. Springer, New York.
- SRIANANTHAKUMAR, S., AND M. L. KING (2006): “A New Approximate Point Optimal Test of a Composite Null Hypothesis,” *Journal of Econometrics*, 130, 101–122.
- STAIGER, D., AND J. H. STOCK (1997): “Instrumental Variables Regression with Weak Instruments,” *Econometrica*, 65, 557–586.
- STOCK, J. H., AND M. W. WATSON (1996): “Confidence Sets in Regressions with Highly Serially Correlated Regressors,” *Working Paper, Harvard University*.
- TANAKA, K. (1996): *Time Series Analysis — Nonstationary and Noninvertible Distribution Theory*. Wiley, New York.

VAN DER VAART, A. W. (1998): *Asymptotic Statistics*. Cambridge University Press, Cambridge, UK.

WALD, A. (1943): “Tests of Statistical Hypotheses Concerning Several Parameters When the Number of Observations is Large,” *Transactions of the American Mathematical Society*, 54, 426–482.